



MYCOTOXINS IN FOODS AND FEEDS 4-FUMONISINS

B.I. Agag

Biochemistry Deficiency Diseases and Toxicology Department,
Animal Health Research Institute, Agricultural Research center

REVIEW ARTICLE

ABSTRACT :

Mycotoxins are toxic metabolites produced by naturally occurring fungi under certain ecological conditions. Mycotoxins contaminated food and feed supplies could increase the economic and health risks to humans and animals. Fumonisin is a group of recently discovered secondary metabolites produced by *Fusarium* spp. especially *F.moniliforme* and occasionally *F.proliferatum* that commonly contaminate corn and corn screenings.

Human consumption of fumonisin-contaminated corn has been associated with increased incidence of esophageal cancer in South Africa, India and China. Fumonisin has been shown to be associated with some major toxicological effects in animals including equine leukoencephalomalacia (ELEM) in horses and porcine pulmonary oedema (PPE) in pigs. Experimentally high doses of fumonisin caused also adverse effects in cattle, sheep, goats, rabbits, poultry and catfish.

Although many domestic animals are susceptible, natural outbreaks of clinical toxicosis from fumonisin have only been reported in horses and swine. Clinical signs of ELEM may include depression, convulsion, ataxia, sweating, apparent blindness, head pressing, recumbency, convulsions and death. The pathognomic lesions are liquefactive necrosis of white matter of brain leaving fluid-filled cavities. Actually affected horses may have centrilobular hepatic necrosis which may be present with or without brain lesions.

Signs of fumonisin toxicosis in swine (PPE) include poor weight gain, weakness, dyspnoea, cyanosis and death. Lesions include pulmonary edema and hydropneumothorax, hepatic and pancreatic necrosis.

INTRODUCTION:

Natural occurrence:

Fumonisin is a group of recently discovered secondary metabolites mainly produced by *Fusarium moniliforme* (*Giberella fujikuroi*) and *F.proliferatum* (Gelderblom *et al.*, 1988; Edrington *et al.*, 1995).

Fumonisin B₁ (FB₁) was isolated in 1988 by Gelderblom *et al.*, and it was chemically characterized by Bezuidenhout *et al.* (1988) and Laurent *et al.* (1989), from cultures of *F.verticillioides* and *F.moniliforme*.

Fumonisin B₁ is produced by other species of *Fusarium* including, *F.anthophilum*, *F.beomiforme*, *F.diamini*, *F.globosum*,

F.hapiforme, *F.nygamai*, *F.oxysporum*, *F.polyphialidicum*, *F.subglutinans* and *F.thapsinum* (WHO, 2000).

The mold is usually found in maize such as screening, that has been damaged by insects or adverse weather conditions (Plumlee, 1997). *F.moniliforme* has also been isolated from pelleted feeds (Plumlee, 1997).

A part from maize and maize products, fumonisins have seldom been found in other products, such as rice (Abbas *et al.*, 1998), asparagus (Logrieco *et al.*, 1998) and sorghum (Shetty and Bhat, 1997). Surveys on other cereals, such as wheat, rye, barley and oats did not show the occurrence of the toxin (Meister *et al.*, 1996).

Maize contaminated naturally by FB₁ can be simultaneously contaminated with other *F.verticillioides* or *F.proliferatum* toxins or with other agriculturally important toxins including deoxynivalenol, zearalenone, aflatoxin and ochratoxin

Many factors including environmental conditions and host susceptibility, determine the incidence and severity of grain mold and subsequent mycotoxin contamination (Stack, 2003). High levels of fumonisins are associated with hot and dry weather, followed by periods of high humidity (Shelby *et al.*, 1994).

Fumonisin levels in raw corn are also influenced by storage conditions and the optimal growth of fumonisin-producing mold that leads to increased levels of fumonisin in the raw corn can occur when the moisture content of harvested raw corn during storage is 18-23% (Bacon and Neslon, 1994). High levels of fumonisins may also occur in raw corn that has been damaged by insects (Bacon and Neslon, 1994; Miller, 1999). The best temperature for the production of FB₁ on corn was 20°C (Weidenborner, 2001).

Types and chemical structure:

The main fumonisins are currently recognized and are designated as fumonisin B₁ (FB₁), fumonisin B₂ (FB₂) and fumonisin B₃ (FB₃), based on the hydroxylation of the hydrocarbon chain (Edrington *et al.*, 1995). FB₁ is the predominating fumonisins in naturally contaminated maize kernels with a ratio of 3:1 (FB₁: FB₂) and 12:1 (FB₁:FB₃) which corresponds to about 70% of the total fumonisins concentration detected. However, in vitro there is some isolates of *F.moniliforme* producing more FB₂ than FB₁ (Weidenborner, 2001).

At least 15 different fumonisins have so far been reported and other minor metabolites have been identified, although most of them have not been shown to occur naturally (WHO, 2000). They have been grouped into four main categories FA₁, FA₂, FA₃ and FAK₁; FB₁, FB₂, FB₃ and FB₄, FC₁, FC₂, FC₃ and FC₄; FP₁, FP₂ and FP₃ (Plattner, 1995; Abbas and Shier, 1997; Musser and Plattner, 1997).

Fumonisin are water soluble, heat stable, alkaline-resistant, aliphatic hydrocarbons with a terminal amine group and two tricarboxylic acid side chains (Osweiler, 1999). The number of position of hydroxyl groups on the aliphatic hydrocarbon determines the structure as FB₁, FB₂ or FB₃ (Bezuidenhout *et al.*, 1988; Steyn, 1995).

The FB₁ form is hydroxylated at the 3, 5 and 10 positions, whereas FB₂ or FB₃ are reduced at the 10 and 5 positions, respectively (Osweiler *et al.*, 1993).

Fumonisin B₁ has the empirical formula C₃₄H₅₉NO₁₅ and is diester of propane-1,2,3-tricarboxylic acid and 2-amino-12,16-dimethyl-3, 5, 10, 14, 15-pentahydroxyicosane (relative

molecular mass:721). The pure structure is a white hygroscopic powder, which is soluble in water, acetonitrile-water or methanol, is stable in acetonitrile-water (1:1), and at food processing temperature and to light (WHO, 2000).

FA₁, FA₂ or FA₃ are the N-acetyl derivatives of FB₁, FB₂ and FB₃, respectively. Within each series different hydroxyl substitution result in different fumonisins FC₁, FC₃ and FC₄ lacking the C-1 terminal methyl group which is characteristic for other fumonisins. In comparison to FC₁, the hydroxylated FC₁(OH-FC₁) has one more hydroxyl group at the C-3 position (Weidenborner, 2001).

Occurrence in foods and feeds:

Maize based human foodstuffs from retail outlet in 5 countries were analysed for FB₁ and FB₂ (Sydenham *et al.*, 1991). The highest man concentration occurred in 2 Egyptian samples 2.38 mg/kg FB₁ and 0.595 mg/kg FB₂. Only of 4 Peruvian samples contained 0.66 mg/kg FB₁ and 0.135 mg/kg FB₂, while only 1 of 2 Canadian samples contained acceptable level of FB₁ (0.05 mg/kg). The corn meal and 10 corn grits products from the USA contained mean concentration of 1.048 mg FB₁ and 0.298 mg/kg FB₂ and 0.601 mg/kg FB₂ and 0.375 mg/kg FB₂, respectively. The mean concentration in 52 corn meal and 15 corn grits samples from South Africa were 0.138 mg/kg FB₁ and 0.083 mg/kg FB₂ and 0.125mg/kg FB₁ and 0.85 mg/kg FB₂, respectively. Only 1 of 10 corn flakes/ lime treated samples contained a low level of FB₁.

Dry milling of maize results in the distribution of fumonisin into the bran, germ and flour (WHO, 2000). In dry milled maize fractions, the fumonisin concentration was approximately 3 times higher in germ and bran than in whole maize, 13 times higher in "C"

flour and 29 times higher than in corn meal and corn grits (Broggi *et al.*, 2002). In experimental wet milling, fumonisin was detected in steep water, gluten fiber and germ, but not in the starch (WHO, 2000).

Fumonisin B₁ levels in animal feedstuffs can be exceptionally high, and reached maximum concentration of 330, 70, 38, 9 and 2 mg/kg in North America, Italy, Brazil, South Africa and Thailand, respectively (WHO, 2000). FB₁ has been recorded to occur in maize and screening servings at levels as high as 195 and 330 mg/kg, respectively (Ross *et al.*, 1991).

The center of veterinary medicine conducted a survey on fumonisin in maize and maize screenings in 1991, in response to an equine leukoencephalomalacia (ELEM) outbreak occurring in late 1990 in the United States. Fumonisin was detected in 13% of the shelled maize samples (1.2 to 3.2 mg/kg), and in 100% of corn screening samples (2.6 to 32 mg/kg) (Price *et al.*, 1993). Another survey found 85% of 158 maize samples (0.6 to 96 mg/kg) (Russel *et al.*, 1993).

In Uruguay all samples of maize-based animal feeds were positive for FB₁. However, highest FB₁ were observed in South Africa for compound feed (11 mg/kg), and Thailand and China for maize (18.8 and 25.97 mg/kg, respectively). In a study of Argentina maize, FB₁ was the major fumonisin at values of up to 11.30 mg/kg (Placinta *et al.*, 1999).

In 1998, random samples of maize and poultry feeds were collected from poultry farms, feed manufacturers and markets in Haryana, India and analyzed for FB₁. 91% of maize samples and 42% of poultry feed samples were found to contain FB₁. FB₁ concentrations in the maize samples ranged from 0.1 to 87 ppm, whereas the poultry feed samples contained FB₁

in the range of 0.2 to 28 ppm (Jindal *et al.*, 1999).

Concentrations of fumonisin in foods and feeds associated with human and animal health problems:

Of several samples obtained from a high oesophageal cancer risk area in the USA 7 of 7 contained FB₁ at levels of 0.105 to 1.915 mg/kg and 6 of 7 FB₂ at levels of 0.07 to 0.46 mg/kg (Sydenham *et al.*, 1991).

The fumonisin B₁ contamination of maize samples collected from two areas of Iran during 1999 was determined (Shephard *et al.*, 2002). The FB₁ levels in Mazandaran province, an area of high esophageal cancer were 0.68-7.66 mg/kg, a mean level of 3.18 mg/kg, while the FB₁ levels found in maize samples from Isfahan, an area of low esophageal cancer were 0.01 to 0.88 mg/kg, a mean level of 0.22 mg/kg.

The fumonisin levels associated with episodes of ELEM and PPE have been found to be higher (Ross *et al.*, 1991). A total of 98 samples of feeds associated with 44 cases of ELEM and 83 samples of feed associated with 44 cases of PPE in USA were analysed for FB₁. Feeds associated with ELEM contained FB₁ ranging from < 1 to 126 mg/kg, with 75 of cases have at least 1 sample > 10 mg/kg. While, feeds associated with PPE ranged from < 1 to 330 mg/kg, with 71% of the cases having at least 1 sample > 10 mg/kg (Ross *et al.*, 1991).

Fourteen feed samples from the USA that were fed to horses prior to the development of LEM were analysed for FB₁ and FB₂ (Thiel *et al.*, 1991 b). All samples contained both FB₁ (1.3-27.0 mg/kg) and FB₂ (0.1-12.6 mg/kg). FB₁ was found to be the major fumonisin in feed samples (53-93%).

A total of 21 *F.moniliforme* contaminated feed sample associated with outbreaks of confirmed and suspected cases of mycotoxicosis

in various animal species (mainly horse and pigs) were collected from farms in the state of Parana, Brazil and analysed for FB₁ and FB₂ (Sydenham *et al.*, 1992). FB₁ and FB₂ were detected in 20 and 18 of the 21 samples, respectively at concentration of 0.2 to 38.5 mg/kg FB₁ and 0.1 to 12.0 mg/kg FB₂.

Absorption, distribution and excretion:

As described by WHO (2000), there is no reports available for fumonisin absorption through inhalation or dermal exposure. However, because fumonisins are present in *F.verticilloides* cells (mycelia, spores and conidiophores) (Tejada-Simon *et al.*, 1995), there is a potential for absorption through inhalation or buccal exposure. The risk from absorption due to animal exposure would seem slight, since fumonisins are very water soluble and, typically polar compounds do not easily penetrate the undamaged skin (Flynn, 1985).

Fumonisin have been reported to be poorly absorbed, rapidly excreted and persistent in small amounts in liver and kidney (Norrd *et al.*, 1996). Fumonisin when dosed orally to vervet monkeys, dairy cows and pigs are poorly absorbed (2 to < 6% of dose) (Prelusky *et al.*, 1994, 1995, 1996 a, b; Shephard *et al.*, 1994a, b).

In ruminants, rumen metabolism may reduce the bioavailability of FB₁ as the hydrolyzed form of FB₁, comprised 60-90% of the total amount of FB₁ found in feces. In non-ruminants the parent compound was the dominant species present (Rice and Ross, 1994). In orally dosed laying hens and dairy cows, the quantity of FB₁ detected in plasma and tissues is very low (< 1% of dose) (Scott *et al.*, 1994; Vudathala *et al.*, 1994 and Prelusky *et al.*, 1996a).

No FB₁ was detected in the milk of lactating sows fed diets containing non-lethal levels of FB₁ and there was no evidence of toxicosis in their suckling pigs (Becker *et al.*, 1995). However, in a study with lactating cows administered FB₁ intravenously, the carry over rate of FB₁ into the milk reached a maximum of 0.11% (Hammer, *et al.*, 1996). In other studies no fumonisins were detected in cows milk (Scott *et al.*, 1994; Richard *et al.*, 1996). FB₁ was found in only one of 165 samples of milk from Wisconsin, USA at a level close to 5 ng/ml (Maragos and Richard, 1994).

There are also no data demonstrating, that fumonisin consumption results in transfer to chicken eggs (Vudathala *et al.*, 1994; Prelusky *et al.*, 1996a). Also, FB₁ was not found in milk, meat or eggs from animals fed grains containing FB₁ levels that would not affect the health of the animals (WHO, 2000).

Mode of action:

Experiments by Wang *et al.* (1991) indicated that fumonisins are inhibitors of sphingolipids biosynthesis through inhibition of ceramide synthetase and by blocking protein synthesis. Sphingolipids are found in large quantities in brain and nerve tissues. Sphingosin is synthesized in the endoplasmic reticulum. Ceramide is formed by a combination of either a free fatty acid or an acyl-coenzyme A and sphingosine (Mayes, 1988).

The site of inhibition occurs where sphingosine and fatty acyl-coenzyme A combine to form dihydroceramide, which would result in accumulation of free sphingosine in tissues and serum (Norred *et al.*, 1992; Merrill *et al.*, 1993 b). It has been postulated that the similarities between fumonisins and long-chain (sphingoid) bases allow them to be recognized as substrate (transition state/or product analogs) of sphingosine or sphinganine-N-acetyl

transferase. Disruption of this pathway could explain at least some of the pathological effects of fumonisins (Leeson *et al.*, 1995).

The degeneration of neuronal cells seen in ELEM may be due to inhibition of sphingolipid biosynthesis because of their high concentration in the brain. On the other hand accumulation of sphinganine in cells exposed to fumonisins may lead to cell death (since long-chain bases are highly cytotoxic), or to cell proliferation, these compounds are mitogenic for some cell types (Wang *et al.*, 1991).

The alteration in sphingolipid metabolism caused by fumonisins can be monitored by measuring serum lipids of sphingosine (So), sphinganine (Sa) and complex sphingolipids (Leeson *et al.*, 1995). Analysis of sphingosine and sphinganine levels in livers and serum revealed an increase in the level of sphinganine, with no change in the level of sphingosine, that result in an increase in the sphinganine: sphingosine ratio (Henery *et al.*, 2000).

Studies in horses (Wang *et al.*, 1991, 1992), pigs (Riley, 1993), sheep (Gurung *et al.*, 1998), broiler chicks (Weibking *et al.*, 1993a), turkeys (Weibking *et al.*, 1993b) and catfish (Goel *et al.*, 1994) have demonstrated a significant increase in the serum Sa to So ratio in animals fed a ration containing FB₁.

Toxicity:

Species difference: FB₁ is extremely toxic to horses, moderately toxic to swine, weakly toxic to cattle and has been associated with oesophageal cancer in humans. Poultry are even more resistant to adverse health effects from fumonisin (Stack, 2003). Pregnant New Zealand rabbits are found to be very sensitive to the toxic effects of FB₁ (LaBord *et al.*, 1997). The species, age and health of the animal as well as the level and duration of exposure to the

mycotoxin will determine the magnitude of the effect of exposure (Stack, 2003).

The feeding of fumonisins contaminated corn produced leukoencephalomalacia (LEM) in horses (Wilson *et al.*, 1990; Ross *et al.*, 1991) and pulmonary edema (PE) and hydrothorax in swine (Harrison *et al.*, 1990; Osweiler *et al.*, 1992). Experimental high doses of fumonisin caused adverse effects in cattle (Osweiler *et al.*, 1993) and poultry (Weibking *et al.*, 1993a). Human consumption of corn has been correlated with increased incidence of oesophageal cancer in certain parts of the world (Sydenham *et al.*, 1991; Chu and Li, 1994). In addition to their adverse effects on the brain, liver and lungs, fumonisins also affect the kidneys, pancreas, testes, thymus, gastrointestinal tract and blood cells (Vencelli and Parker, 1999).

Fumonisin have produced liver damage and changes in the levels of certain classes of lipids especially sphingolipids, in all animal studies (Merill *et al.*, 1997), kidney lesions were also found in many animals (Norred *et al.*, 1998; Merrill *et al.*, 1997).

Chronic feeding of purified FB₁ at levels of 50 ppm or more produced liver cancer and decreased life span in female mice and kidney cancer in male rats without decrease life spans (NTP, 1999).

Disturbances of thermoregulation has been observed in some animals exposed to the mycotoxin (Becker *et al.*, 1995). In horses, FB₁ causes profuse sweating, the major cooling mechanism for this species, even though body temperature remain normal (McCue, 1989). Cattle with fumonisin toxicosis has hyperthermia, with heat intolerance developing at a lower body temperature than normal for cattle (Schmidt and Osborn, 1993; Thompson and Stuedemann, 1993). Although,

hyperthermia has not been reported in swine with FB₁ toxicosis, the reduced lung function associated with edema would reduce the ability to pant. In swine panting is a major mechanism for coping with high temperature, thus high environmental temperature may exacerbate FB₁ toxicosis attributable to reduced pulmonary function associated with FB₁ induced pulmonary edema (Becker *et al.*, 1995).

Human health effects:

Currently, there is no direct evidence that fumonisins cause adverse health effects in humans. Studies currently available demonstrate only inconclusive association between fumonisins and human cancer (FDA, 2001b).

A very high incidence of oesophageal cancer among the black population of the Transkei, South Africa has been reported in several surveys (Jaskiewicz *et al.*, 1987; Makaula *et al.*, 1996). Investigation in South Africa suggested an association between high levels of fumonisin producing molds on corn used to make alcoholic beverages and oesophageal cancer in humans (Rheeder *et al.*, 1992). This corn has been found to contain up to 118 mg/kg fumonisins.

Based on experiments conducted on beer made from wort containing FB₁, such beers could contain fumonisin concentration of 30 mg/liter beer (Scott *et al.*, 1995). However, these studies were limited by the lack of controlled conditions, particularly for established confounding risk factors (e.g. alcohol consumption) and therefore do not allow any definite conclusions to be made about cancer causation in humans (FDA, 2001 b).

An outbreak of poisoning, characterized by abdominal pain and diarrhea caused by fumonisin-contaminated maize and sorghum in India during 1995, was reported (Bhat *et al.*,

1997). An epidemiological survey was conducted in the affected villages and a detailed house to house in selected villages. People in 27 out of 50 villages surveyed in Karnataka state were affected and disease was seen only in households and subjects consuming rain damaged moldy sorghum or maize.

All 20 sorghum and 12 maize samples collected from household had fusarium sp. as the dominant microflora and contained FB₁ in the range of 0.14 to 7.8 and 0.25 to 64.7 mg/kg, respectively. In contrast, samples collected from unaffected households has FB₁ in low levels ranging from 0.07 to 0.36 and 0.05 to 0.24 mg/kg, respectively.

However, this study lacked control of established risk factors. In addition, contaminants other than mycotoxins can not be eliminated as causative factors, and a similar dissociation was not detected in studies conducted in other countries (FDA, 2001 b).

Other studies associated with high levels of fumonisin producing molds on corn with oesophageal cancer in China reported that corn samples from areas with high incidence of esophageal cancer, contained FB₁ at levels ranging from 18 to 155 mg/kg. These results established that home-grown maize in high incidence areas of oesophageal cancer in China may be contaminated with very high levels of FB₁. These studies were correlation studies where there was no clear picture on the association of either fumonisins or *F.verticillioioides* contamination with oesophageal cancer (WHO, 2000).

Further in an area of China with high incidence of gastric cancer, Groves *et al.* (1999) observed a lack of association between consumption of fumonisin-contaminated corn with gastric or any other human cancer.

Animal health effects:

Ruminants:

Cattle and sheep are mildly affected by dietary concentrations of fumonisin greater than 100 ppm, but fatalities do not occur (Osweiler *et al.*, 1993). Very few studies have focused on the effects of fumonisins in ruminants. (Kriek *et al.*, 1981) demonstrated that *F.moniliforme* cultures capable of inducing ELEM and PPE were also toxic to sheep. Beasley and Buck (1982) reported feed refusal in cattle given corn contaminated with *F.moniliforme*, but existence of fumonisins was not known at that time.

Dietary fumonisins concentration greater than 148 mg/kg of diet to calves may adversely affect liver and immune functions (Osweiler *et al.*, 1993). Livers from treated calves had mild hydropic degeneration and cloudy swelling in a peri-acinar pattern throughout the liver. These changes were consistent with the mild to moderate elevation of liver enzymes. Cattle fed 530 mg of total fumonisin /kg diet, supplied by fumonisin-containing culture material for 30 days, showed no signs of toxicosis (Smith and Thakur, 1996), although liver function tests revealed some hepatobiliary compromise.

Fumonisin B₁-contaminated feed (95 mg/kg), fed for 112 days caused no treatment-related effects on feed intake, nutrient digestibility, or weight gain in weanling angora goats (Gurung *et al.*, 1998). However, sphingolipid disruption and mild liver and kidney damage were observed in weanling goats.

In contrast, (Edrington *et al.*, 1995) observed clinical signs such as marked decrease in feed intake diarrhea, lethargy and ultimately death in lambs dosed intraruminally with 11.1,

22.2 and 45.5 mg of total fumonisins/ kg of b.wt. for only 4 days. The actual amounts of total fumonisin intake were determined to be 355.2, 710.4 and 1456 mg/ lamb/ day. Such as concentrations have not been reported to occur under normal feeding conditions (Gurung *et al.*, 1998).

Monogastric animals:

Equidae:

Horse are the most sensitive of the domestic animals and dietary concentration as low as 10 ppm for more than 50 days may cause leukoencephalomalacia (LEM) (Wilson *et al.*, 1990; Ross *et al.*, 1992). The ELEM has been recognized since 19th century as a sporadically occurring condition (WHO, 2000), and was experimentally produced by feeding moldy maize obtained from a field case in Kansas by Butler in 1902. The disease was known as "moldy corn poisoning" but attempts to identify the responsible fungus failed. History has referred to this disease by many names including: moldy corn poisoning, corn stalk disease, forage poisoning, leukoencephalomyelitis, blind staggers, epizootic cerebritis and non virus encephalomyelitis (Wilson *et al.*, 1985; Asquith, 1991; Masri, 1992).

As described by WHO (2000), Wilson and Maronport (1971) succeeded in establishing the causative agent when they isolated *F.verticillioides* as the predominant contaminant of moldy maize that has caused cases of ELEM in Egypt, and reproduced the disease by feeding culture material of the fungus on maize to 2 donkeys. Subsequently investigators in South Africa confirmed the ability of *F.verticillioides* (MRC826) culture material to induce the characteristic clinical signs and pathological changes of ELEM as well as hepatosis in horses

and donkeys (Kellerman *et al.*, 1972; Marasas *et al.*, 1976, 1988 and Kriek *et al.*, 1981).

ELEM has also produced in horse by intravenous administration of pure FB₁ (Bezuidenhout *et al.*, 1988; Gelderblam *et al.*, 1988) and in horses given pure FB₁ by stomach tube (Kellerman *et al.*, 1990). Moreover the disease has been observed naturally in horses and ponies fed feeds contaminated with fumonisins (Wilson *et al.*, 1992; Ross *et al.*, 1993).

The lowest FB₁ dose that has resulted in ELEM in a controlled experiment is 22 mg/kg in diets formulated with naturally contaminated maize screenings (Wilson *et al.*, 1992). Ross *et al.* (1992) suggested that concentrations ≥ 10 mg/kg FB₁ in horse feeds were likely to be involved in LEM. In 40 to 45 confirmed cases of ELEM, FB₁ levels in feeds were ≥ 10 mg/kg, while non-problem feeds contained ≤ 10 mg/kg FB₁.

Clinical signs:

The clinical signs of ELEM appear at a variable length of time after feeding of the mycotoxin is begun (Kibluk *et al.*, 1995). In experimental cases, clinical signs have appeared as early as day 7 and as late as day 75 (Ross *et al.*, 1993). Clinical signs of the disease usually appear abruptly and the clinical course is relatively short, with most affected horses dying within 2 to 3 days (Wilson *et al.*, 1985; Marsi *et al.*, 1987; Kobuk *et al.*, 1995).

The first sign may be inappetence, followed quickly by neurologic abnormalities. Behavioral signs vary from depression and a pathy to apprehension, hyperexcitability or frenzy (Kellerman *et al.*, 1990; Masri *et al.*, 1992; Ross *et al.*, 1993; Kobluk *et al.*, 1995). Other neurologic signs may include head pressing, aimless circling, unilateral or bilateral blindness, ataxia, tremors, cranial nerve deficits

(such as lip and tongue paresis and dysphagia), and weakness, progression of the condition leads to recumbency, convulsions, coma and death (Wilson *et al.*, 1985, 1990; Masri *et al.*, 1987; Marasas *et al.*, 1988; Kellerman *et al.*, 1990; Thiel *et al.*, 1991b; Ross *et al.*, 1993 and Konluk *et al.*, 1995). In a horse injected intravenously 7 times over 10 days with 0.125 mg of FB₁/kg b.wt./day, clinical signs of neurotoxicosis included nervousness followed by a pathy, a wide-based stance, trembling, ataxia, reluctance to move, paresis of the lower limb and tongue and an inability to eat or drink were recorded (Marasas *et al.*, 1988). The principal lesions were severe oedema of the brain and early bilaterally symmetrical focal necrosis in the medulla oblongata.

Death frequently occurs within 5 days of the onset of clinical signs (Smith, 2000): sudden death, without prior clinical signs, has been reported. Animals that survive often have persistent neurologic deficit (Plumlee, 1997). The hepatic form of the disease may be accompanied by icterus, oedema, hemorrhage and hepatoenkephalopathy (Plumlee, 1997).

Gross and histopathological findings: ELEM syndrome is characterized macroscopically by the presence of liquefactive necrotic lesions in the white matter of the cerebrum (WHO, 2000). Other lesions include encephalomalacia of both white and grey matter in the cerebrum, cerebellum, midbrain or brain stem (Masri *et al.*, 1987; Konluk *et al.*, 1995). Acutely affected horses may have centrilobular hepatic necrosis, which may be present with or without the brain lesions (Marasas *et al.*, 1988; Wilson *et al.*, 1990).

Microscopic lesions included: liquefactive necrosis of white matter, with complete loss in some areas perivascular edema, hemorrhage

and perivascular cuffing in the white matter adjacent to malacic foci. Eosinophils and plasma cells predominated in perivascular cuffs. Foci of myelin degeneration were seen in non-necrotic areas of white matter. Lesions in grey matter included satellitosis and neuronophagia. Leptomeningeal edema was present in some cases (Marasas *et al.*, 1976; Rooney and Robertson, 1996).

In conjunction with ELEM, muscular melting and cardiac failure have been reported (Vencilli and Parker, 1999). In addition to brain lesions, histopathological abnormalities in liver and kidney have been reported in horses orally dosed with pure fumonisins, maize screening, naturally contaminated with fumonisins, or culture material containing known amounts of fumonisins (Kellerman *et al.*, 1990; Wilson *et al.*, 1992; Caramalli *et al.*, 1993 and Ross *et al.*, 1993). Lesions of hepatic form include centrilobular necrosis, periportal fibrosis, periportal vacuolation and bile duct proliferation; lipofuscin deposition in macrophages and Kupffer cells, and infiltration with a mixture of inflammatory cells (Marasas *et al.*, 1976; Plumlee, 1997). Perirenal edema and congestion may be present (Marasas *et al.*, 1976).

Elevated serum enzyme level indication of liver damage (Wilson *et al.*, 1992) are produced by elevation of the serum Sa/So ratio (Wang *et al.*, 1992). The protein concentration and white cell count in the cerebrospinal fluid are elevated (Plumlee, 1997).

Natural occurrence of ELEM:

Outbreak of ELEM has been described throughout the world in Argentina, China, Egypt, New Caledonia, South Africa, Greece and other parts of Europe and the United States

(Jubb and Kennedy, 1985 and Marasas *et al.*, 1988).

The disease occurs most commonly in the United States from September to May. Certain growing and harvest conditions (dry summers and/or drought, followed by rainy autumns) increase the amount of *Fusarium moniliforme* on feed corn (Rooney and Robertson, 1996). Epizootics have occurred in years in which corn was harvested during periods of high humidity (Badiali *et al.*, 1968; Masri *et al.*, 1987).

In 1934 to 1935, an episode of ELEM was recorded in Illinois (Graham, 1936), where 5000 horses died following a summer drought and early fall rains, when horses were allowed to forage in the stalk fields, and others were fed moldy corn after harvesting. Sporadic occurrences of the disease were reported in Egypt by Badiali *et al.* (1968), following the flooding of the Nile, resulting in mold growth on the autumn crop which was fed to donkeys.

An outbreak of ELEM was reported in a riding club in New Caledonia in 1981, where horses were being fed a ration containing high levels of locally grown corn. The development of the disease coincided with the feeding of a new batch of feed containing the corn. Out of the 45 affected horses, 5 died 10 to 28 days after the onset of clinical signs. *F.moniliforme* was recovered from all samples of corn that were examined. A second episode occurred in 1983 involving a 15-year-old stallion that had been fed locally produced corn. In this case, the major gross pathology was a focus of liquefaction of the white matter in the right cerebral hemisphere (Domench *et al.*, 1985; Asquith, 1991).

A survey conducted in 1984 to 1985, evaluated 22 farms with confirmed cases of ELEM in North Carolina. Gross lesions of necropsied animals revealed liquefaction

necrosis in the white matter of one or both cerebral hemispheres. Preliminary data suggested that some of the oats used in the feed preparation may have been infected. This was the first report of the disease occurring from the ingestion of commercial pelleted and non-pelleted horse rations and indicated that *F.moniliforme* may be capable of remaining viable even after the pelleting process (Wilson *et al.*, 1985; Asquith, 1991).

During the autumn of 1989, in an ELEM involving 18 of 66 purebred Arabian horses at breeding, training stable in Arizona, USA, the condition was fatal in 14 horses (Wilson *et al.*, 1990). These horses had been fed a diet containing 37 to 122 ppm of FB₁. Gross pathological findings included, liquefactive necrosis in parts of the cerebral white matter and hemorrhagic foci of various sizes in the brain stem. Histopathological findings included reified white matter with pyknotic nuclei and eosinophilic cytoplasm. This was the first definite report on ELEM and associated FB₁ concentration.

An outbreak of ELEM affecting 6 of 10 pleasure horses in adjacent paddocks at a boarding facility in Pennsylvania, USA is reported (Wilkins *et al.*, 1994). The horses showed depression, lethargy, ataxia, blindness, seizures and recumbency after eating whole ear maize. Four horses died or were euthanized. *F.moniliforme* (*Gibberella fujikuroi*) was cultured from the ear maize and fumonisins B₁, B₂ and B₃ at concentrations of 370, 105 and 41 ppm, respectively, were detected.

The first report of ELEM in Italy was observed by Caramelli *et al.* (1993) on a farm in Piedmont, where 3 of 40 horses showed sudden anorexia, hyperexcitability, incoordination, flacid lips, blindness and recumbency followed by death in 72 hours. These horses had been fed

for some days on maize screenings containing 60 mg/kg FB₁ and 14.6 mg/kg FB₂.

ELEM was first diagnosed in Hungary in September 1995, in a 13- years- old castrated male, a 6- ears- old stallion and a 2- years- old foal (Fazekas and Bajmicy, 1996). The disease started with lack of appetite followed by difficulties in swallowing and chewing indicating the paralysis of cephalic and pharyngeal muscles. Paralysis of the cephalic and cervical muscles spread to the muscles of the extremities and trunk and ataxia developed. Signs of blindness appeared in an animal. At the final stages of the disease, the affected animals became recumbent and died. Post-mortem examination of one horse showed oedema of the lungs and white matter of the brain. The affected animals were fed green maize containing FB₁ at 18.5 mg/kg of the whole plant.

Pigs:

Swine consuming more than approximately 120 ppm dietary fumonisins for 4 to 10 days develop acute porcine pulmonary oedema (PPE) (Coloin and Harrison, 1992; Colvin *et al.*, 1993), more than 50 ppm of fumonisins causes mild liver lesions within 7 to 10 days, but concentrations of 25 ppm or less cause no apparent clinical effects. Mild microscopic hepatic lesions can be produced from diets as low as 23 ppm (Mortelin *et al.*, 1994; Osweiler, 1999). The sphinganine- sphingosine ratio may be altered in the serum by feed concentrations as low as 5 ppm and in the lung, liver and kidney by feed concentrations of 10 ppm, although the clinical significance of this is not established (Mortelin *et al.*, 1994; Riley *et al.*, 1993 and Rotter *et al.*, 1996).

PPE represents the acute effects of FB₁ (Becker *et al.*, 1995), while, the chronic effects of FB₁ in weaned pigs showed hyperplastic plaques

in the oesophageal mucosa (Casteel *et al.*, 1993). Swine fed a ration containing 100- 190 ppm FB₁ for up to 83 days developed nodular hyperplasia of the liver and hyperplasia of the esophageal mucosa (Casteel *et al.*, 1993). Swine fed 150- 170 ppm FB₁ for up to 210 days developed medial hyperplasia of the pulmonary arteries and changes in gravimetric measurements of the heart (Casteel *et al.*, 1994). In experimental trials, culture material of *F.moniliforme* (NRC826) was fed to horses, pigs, sheep, rats and baboons (Krick *et al.*, 1981) lung oedema occurred only in pigs.

Acute PPE might be induced by increases in pulmonary interstitial macrophages resulting in release of vasoactive mediators (Huschek *et al.*, 1992), pulmonary hypertension caused by vasoconstriction (Smith *et al.*, 1996) and acute left- sided heart failure (Smith *et al.*, 1999). There responses have been hypothesized to lead to pulmonary oedema either by increased pulmonary hydrostatic pressure or by pulmonary capillary endothelial cell damage (Cumpecht *et al.*, 1998; Osweiler, 1999).

Clinical signs of PPE typically occur soon (2-7 days) after pigs consume diets containing large concentrations of fumonisins over a short period of time (WHO, 2000). These signs include labored or difficult breathing (dyspnoea), weakness, cyanosis and death (Osweiler *et al.*, 1992; Mortelin *et al.*, 1994 and Stack, 2003). Once signs appear, death may occur in less than 4 hours (Colvin and Harrison, 1992; Riley *et al.*, 1993). Pregnant sows may abort from 1 to 4 days after the onset of signs (Osweiler, 1992), probably as a sequela to fetal anoxia from severe pulmonary oedema (Osweiler, 1999).

At necropsy, the diseased animals exhibit varying degrees of interstitial or interlobular oedema, with pulmonary oedema and

hydrothorax. Varying amounts of clear cell-free yellow fluid accumulate in the pleural cavity (Colvin and Harrison, 1992; Colvin *et al.*, 1993). Early histologic changes in the lung consisted of perivascular oedema followed by interlobular and peribronchial oedema. Ultrastructurally, alveolar endothelial cells contained unique accumulations of membranous material in the cytocavity network (Cumpecht *et al.*, 1998). Acidophilic fibrillar material in alveoli and interlobular lymphatics, hyalinized alveolar capillary thrombi and increased numbers of pulmonary intravascular macrophages filled with osmophilic material (Colvin *et al.*, 1992; Riley *et al.*, 1993 and Osweiler, 1999).

Dietary fumonisin concentrations of 75 to 120 ppm for 1 to 4 weeks cause icterus, reduced feed intake weight loss and occasionally diarrhoea (Osweiler *et al.*, 1993; Osweiler, 1999). Toxic hepatitis occurs concurrently with PPE (Osweiler *et al.*, 1992; Colvin *et al.*, 1993) and is also observed in animals that consume high levels of fumonisins but do not develop PPE (Hascckek *et al.*, 1996). Typically the liver contains multiple foci of coagulative necrosis with disrupted hepatic architecture, increased mitotic figures in hepatocytes and single hepatic necrosis (Osweiler *et al.*, 1992; Colvin and Harrison, 1992; Colvin *et al.*, 1993 and Riley *et al.*, 1993).

Rabbits and Poultry:

Rabbits:

Rabbits given multiple doses of FB₁ (0.15, 0.3, 0.5 or 1.0 mg/kg b.wt. for 4 or 5 days) intravenously, were lethargic and anorectic and had decreased urine production (Gumprecht *et al.*, 1995). Renal lesions consisted of severe proximal tubular necrosis. Liver lesions were variable and consisted of mild hepatic vacuolation and bile stasis. A single dose of FB₁

at 1.0 mg/kg b.wt induced renal but not hepatic injury.

Bucci *et al.* (1996) studied LEM and hemorrhage in the brain of rabbits gavaged with FB₁ (1.75 mg/kg b.wt. daily) and observed that of 5 gavaged pregnant rabbits, 2 died after receiving 9 and 13 doses, respectively. Microscopic examination revealed focal small hemorrhages in cerebral white matter in both animals, with malacia and hemorrhage also present in the hippocampus of one. Both animals also had marked degeneration of renal tubules epithelium and of hepatocytes. Apoptosis was the dominant degenerative change in kidney and liver.

Pregnant New Zealand white rabbits are very sensitive to the toxic effects of Fb₁ (LaBorde *et al.*, 1997). Maternal toxicity was observed at daily gavage dosage 0.25 mg/kg b.wt from gestational day 3 to gestational day 19. The maternal kidney, serum and urine Sa/So ratios were increased, but there was no increase in these ratios in fetal liver, brain or kidney. The lowest observed effect level for maternal toxicity was 0.1 mg FB₁/kg b.wt., which is equivalent to a calculated dietary FB₁ level of 2.3 mg/kg diet (LaBorde *et al.*, 1997; WHO, 2000). This level is very close to the 2.24 mg/kg of fumonisin in the diet of young colts that could be theoretically be hazardous. Consequently, rabbits were grouped with horses (FDA, 2001a).

Poultry:

Turkey poults are more sensitive to the toxic effect of FB₁ than broiler chicks (Weibking *et al.*, 1993b) but chickens, turkeys and ducklings are relatively resistant to FB₁ in comparison to horses and swine, where significant toxicity and mortality occur at levels greater than 10 and 20 mg FB₁/ kg diet, respectively (Ross *et al.*, 1991;

Oswailer *et al.*, 1992). On the other hand, one day- old chicks are more sensitive to the adverse effects of FB₁ than 7 or 21 days- old chicken (Javed *et al.*, 1993).

Broilers:

A day- old chicks fed diets containing 200 or 300 mg FB₁/kg diet for 3 weeks had thymic cortical atrophy, multifocal hepatic necrosis, biliary hyperplasia and widening of the proliferating cartilage zone in proximal tibiotarsal physes (Ledoux *et al.*, 1992). Acute randomly oriented hepatic necrosis with moderate biliary hyperplasia and hypertrophy of kupffer cells, mild- villus atrophy and goblet cells hyperplasia in the lower small intestines, small foci of acute myocardial and skeletal muscle necrosis and widening of both proliferating and hypertrophic zones in proximal tibiotarsal physes were found in a day- old chicks fed 300mg FB₁/kg diet for 2 weeks (Brown *et al.*, 1992).

A day- old chicks fed diets containing 89, 190, 283, 289, 481, 592 and 681 mg of total fumonisins/ kg diet for 3 weeks, had increased sphinganine: sphingosine ratios. Isolated foci of hepatic necrosis with a mild heterophil and macrophage infiltration, moderate diffuse hepatocellular hyperplasia, mild biliary hyperplasia and moderate to severe periportal granulocytic cell proliferation were noted only in broiler fed levels equal to or greater than 289 mg/kg (Weibking *et al.*, 1993a; FDA, 2001a).

Dose response clinical signs, reduced body weight and mortality were observed in chicken fed diets containing 125 or 274 mg purified FB₁/kg diet (Javed *et al.*, 1993).

Immunosuppression in chickens was produced in broilers fed diets containing 10 mg pure FB₁/kg diet or diets formulated from *F.verticillioides* (MRC826) culture material

containing 30 to 80 mg FB₁/kg diet. Birds had reduced spleen and/ or bursa weights and altered haematological parameters (Espada *et al.*, 1994, 1997).

Decreased prothrombin time and increased fibrinogen concentration and the activity of antithrombin III were observed in blood from broiler chicks that had been fed dietary FB₁ at a concentration of 10 mg/kg for only 6 days (Henry *et al.*, 2000). Dietary FB₁ at concentration of 80 mg/kg or less did not adversely affect body weight, feed efficiency or water consumption in broiler chicks. Liver sphinganine concentration and the sphinganine: sphingosine ratio were increased significantly in all treated groups that had received 20, 40 and 80 mg FB₁/kg diet (Henry *et al.*, 2000).

Ducks:

A day- old white Pekiny ducklings fed diets containing 120.5, 240.9 and 481.8 mg of total fumonisins for 21 days had statistical increase in the mean kidney weights, liver sphinganine: sphingosine ratio and serum gamma glutamyl transferase activity as well as, mild to moderate hepatocellular hyperplasia (Bermudez *et al.*, 1995; FDA, 2001a).

Turkeys:

Turkey poults have been shown to be susceptible to the adverse effects of FB₁. Biliary hyperplasia, hypertrophy of kupffer cells, thymic cortical atrophy and moderate widening of the proliferating and degenerating hypertrophied zones of tibial physes were observed in turkey poults fed diets containing 100-200 mg FB₁/ kg diets (Weibking *et al.*, 1993b). Poor performance, increased organ weight, diarrhea, biliary hyperplasia, hepatocellular hyperplasia and rickets are

caused in poult by dietary levels of FB₁ (≥ 75 mg/kg) (Weibking *et al.*, 1993a, 1995).

A day- old turkey poult fed diets containing graded levels of total fumonisins (33, 66, 99, 132, 231, 330, 429, 528 and 627 mg/kg) for 21 days showed liver lesions at levels equal to or greater than 99 mg/kg. hepatocellular hyperplasia was mild at 99 and 132 mg/kg, moderate to severe at 330 mg/kg and severe at 429, 528 and 627 mg/kg (Ledoux *et al.*, 1996; FDA, 2001a).

Turkey poult fed FB₁- contaminated diet for 21 days showed mild hepatic lesions at 75 mg/kg, cardiac lesions at 475 mg/kg, decreased performance at 325 mg/kg and increased liver weights at 25 mg/kg (Ledoux *et al.*, 1996). Increased liver weight with moderate diffuse hepatocellular hyperplasia and elevated enzymes were observed in poult fed 200 mg FB₁/ Kg of diet (Bermudez *et al.*, 1997).

Gross lesions induced by fumonisin toxicity in poultry comprise consistent enlargement of the liver, and variable enlargement of kidney, pancreas, proventriculus and gizzard, atrophy of lymphoid organs and rickets (Calnek *et al.*, 1997).

Histologically, the liver has multifocal necrosis of hepatocytes, hyperplasia of hepatocytes and bile ductules and hypertrophy of Kupffer cells. The intestine has villus atrophy and goblet cell hyperplasia. Growth plates are widened in both the zones of proliferating and hypertrophic cartilage. Myocardium and skeletal muscles have mild lesions and lymphoid tissues are depleted (Gureshi and Hagler, 1992; Calnek *et al.*, 1997).

From the results of experimental studies conducted on fumonisins in poultry, it appears that: the turkey poult study (Ledoux *et al.*, 1996) showed toxic effects at 99 ppm, the broiler study (Weibking *et al.*, 1993a) at 190 to 280 ppm and the duckling study (Bermudez *et al.*, 1995) at

120 ppm of total fumonisins in the diet. These toxic effects center around liver lesions. although chicken may be slightly more resistant to fumonisins than turkeys and duckling, it appears that these 3 species should be considered fairly resistant to the toxic effects of fumonisins and should be grouped into one category (poultry fed for slaughter) (FDA, 2001a).

Catfish:

In channel catfish dosed with *F.moniliforme* culture, the lowest dietary concentrations of fumonisin associated with significantly elevated ratios of sphinganine to sphingosine in kidney, serum, liver and muscle were 10, 20, 40 and 80 mg of FB₁/ kg of diet, respectively (Goel *et al.*, 1994). Catfish fed diets containing FB₁ levels equal to or higher than 40 mg/kg of diet for 12 weeks showed decreased feed consumption, feed efficiency and weight gain (Li *et al.*, 1994; Li and Robinson, 1995). The minimum level of FB₁ that depressed growth was 20- 40 mg/kg (Li *et al.*, 1994).

Catfish fed diets containing 40 mg FB₁/ kg diet for 12 weeks had increased liver glycogen, increased vacuolation in nerve fibers and perivascular lymphohistiocytic investment in the brain compared with fish fed diets containing lower levels of fumonisins (0.7 to 20 mg/kg) (Li *et al.*, 1994). Small white foci (2 to 4 mm diameter) of subcapsular adipocyte hyperplasia were observed in the livers of channel catfish fed 20 mg of FB₁/ kg of diet for 10 and 14 weeks. Livers had also swollen hepatocytes with lipido containing vacuoles, lymphocyte infiltration and scattered necrotic hepatocytes (Lumlertdacha *et al.*, 1995).

Regulatory control:

Based on the wealth of available information on the adverse animal health effects associated with fumonisins, human health risks associated with exposure to fumonisins are possible. Therefore, human exposure to such natural toxins should not exceed levels achievable with the use of good agricultural and good manufacturing practices (FDA, 2001b).

The recommended maximum levels for fumonisins in human foods and animal feeds (Tables, 1 and 2) that FDA considers achievable with the use of good agricultural and good manufacturing practices are presented below (FDA, 2000; FDA, 2001 b, c).

Table (1): Recommended maximum levels of fumonisins in human food products, U.S.

Product	Total fumonisins (FB ₁ + FB ₂ + FB ₃)
Degermed dry milled corn products (e.g., flaking grits, corn grits, corn meal, corn flour with fat content of < 2.25%, dry weight basis)	2
Whole or partially degermed dry milled corn products (e.g. flaking grits, corn grits, corn meal, corn flour with content of < 2.25%, dry weight basis)	4
Dry milled corn bran	4
Cleaned corn intended for masa production	4
Cleaned corn intended for popcorn	3

Table (2) Recommended maximum levels for fumonisins in corn, corn by products and the total ration for various animals species

Animal or class	Maximum level of total fumonisins in corn and corn products (ppm ¹)	Feed factor ²	Maximum levels of total fumonisins in total ration (ppm) ¹
Horses ³	5	0.2	1
Rabbit	5	0.2	1
Catfish	20	0.5	10
Swine	20	0.5	10
Ruminants ⁴	60	0.5	30
Milk ⁵	60	0.5	30
Poultry ⁶	100	0.5	50
Ruminant, poultry and mink breeding sock ⁷	30	0.5	15
All others ⁸	10	0.5	5

1. Total fumonisins FB₁ + FB₂ + FB₃.
2. Fraction of corn or corn by- products mixed into the total ration
3. Includes asses, zebra and onagers.
4. Cattle, sheep, goats and other ruminants that are ≥ 3 months old and fed for slaughter.
5. Fed for pelt production.
6. Turkeys, chickens, ducklings and other poultry fed for slaughter.
7. Includes laying hens, roosters, lactating dairy cows and bulls.
8. Includes dogs and cats.

In six instances, the U.S. FDA (2000) grouped species together because the animals seemed to have a similar sensitivity to fumonisins. This is an attempt to avoid a multitude of guidance levels and does not necessarily imply that the species are biologically similar.

1-Horses and rabbits were grouped together as the most sensitive species. Corn and corn- by

products used in rations of horses and rabbits should contain less than 5 ppm of total fumonisins and comprise no more than 20% the dry weight of the total ration (Table 2). The total ration should contain less than 1 ppm of total fumonisins.

2-Catfish and swine were grouped together as intermediate in sensitivity to fumonisins. Corn and by products used in rations of catfish and

swine should contain less than 20 ppm of total fumonisins, and comprise no more than 50% of the dry weight of the total ration. The total ration should contain less than 10 ppm of total fumonisins.

3-Ruminants, mink and poultry were considered more resistant than horses, rabbits, catfish and swine to fumonisins. Corn and corn by products used in rations of ruminants that are at least 3 months old fed for slaughter and in rations of mink fed for pelt production should contain less than 60 ppm of total fumonisins, and comprise no more than 50% of the dry weight of the total ration. The total ration should contain less than 30 ppm of total fumonisins.

4-Corn and corn- by products used in the rations of poultry fed for slaughter should contain less than 100 ppm of total fumonisins, and comprise no more than 50% of the dry weight of the total ration. The total ration should contain less than 50 ppm of total fumonisins.

5-Corn and corn- by products used in the rations of mink, ruminants and poultry breeding stock should contain less than 30 ppm of total fumonisins, and comprise no more than 50% of the weight of the total ration. The total ration should contain less than 15 ppm of total fumonisins.

6-Corn and corn- by products used in rations of dogs and cats should contain less than 10 ppm of total fumonisins, and comprise no more than 50% of the dry weight of the total ration. The total ration should contain less than 5 ppm of total fumonisins.

REFERENCES :

- Abbas, H.K. and Shier, W.T. (1997): "Phytotoxicity of australifungin and fumonisins to weeds. In: proc. of the 1997 Brighton Crop Protection Conference-Weeds." Croydon, UK, British Crop Protection Council, Pp. 795- 800.
- Abbas, H.K.; Cartwright, R.D.; Shier, W.T.; Abouzied, M.M.; Bird, C.B.; Rice, L.G.; Ross, P.F.; Sciumbato, G.L. and Meredith, F.I. (1998): "Natural occurrence of fumonisins in rice with fusarium sheath rot disease." *Plant Dis.*, 82: 22- 25.
- Asquith, R.L. (1991): "Mycotoxicosis in horses" In: Smith, J.E. and Hederson, R.S. (eds.) *mycotoxicosis and animals foods*. PP. 679- 687- CRC Press, Inc., London.
- Bacon, C.W. and Nelson, P.E. (1994): "Fumonisin production in corn by toxigenic strains of *Fusarium moniliforme* and *Fusarium proliferatum*." *J. Food Protection*, 57 (6): 514- 521.
- Badiali, L.; Abou- Youssef, M.H.; Rodwan, A.L.; Hamdy, F.M. and Hildebrandt, P.K. (1968): "Moldy corn poisoning as the major cause of an encephalomalacia syndrome in Egyptian equidae." *Am. J. Vet. Res.*, 29: 2020.
- Beasley, V.R. and Buck, W.B. (1982): "Feed refusal in cattle associated with *Fusarium moniliforme* in corn." *Vet. Rec.*, 111: 393.
- Becker, B.A.; Pace, L.; Rottinghaus, G.E.; Shalby, R.; Misfeldt, M. and Ross, M.S. (1995): "Effects of feeding fumonisin B₁ in lactating sows and their sucking pigs." *Am. J. Vet. Re.*, 56 (9): 1253- 1258.
- Bermudez, A.J.; Ledoux, D.R. and Rottinghaus, G.E. (1995): "Effects of *Fusarium moniliforme* culture material containing known levels of fumonisin B₁ in ducklings" *Avian Dis.*, 39 (4): 879- 886.
- Bermudez, A.J.; Ledoux, D.R.; Rottinghaus, G.E. and Bennett, G.A. (1997): "The individual and combined effects of the

- fusarium mycotoxins moniliformin and fumonisin B₁ in turkeys." *Avian Dis.*, 41: 304- 311.
- Bezuidenhoudt, S.C.; Wentzel, A.; Gelderblom, W.C.A. (1988): "Structure elucidation of the fumonisins, mycotoxins from *Fusarium moniliforme*." *J. Chem. Soc. Chem. Commun.*, 743- 745.
- Bhat, R.V.; Shetty, P.H.; Amruth, R.P. and Sudershan, R.V. (1997): "A foodborne disease outbreak due to the consumption of moldy Sorghum and maize containing fumonisin mycotoxins" *Clin. Toxicol.*, 35: 249- 255.
- Broggi, L.E.; Resnik, S.L.; Pacin, A.M.; Gonzalez, H.H.; Cano, G. and Taglieri, D. (2002): "Distibuion of fumonisins in dry milled corn fractions in Argentina." *Food Assitives and ontaminants*, 19 (5): 465- 469.
- Brown, T.B.; Rottinghaus, G.E. and Williams, M.E. (1992): "Fumonisin mycotoxicosis in broilers: performance and pathology." *Avian Dis.*, 36: 450-454.
- Bucci, T.J.; Hansen, D.K. and LaBorde, J.B. (1996): "Leukoencephalomalacia and hemorrhage in the brain of rabbits gavaged with mycotoxin fumonisin B₁." *Natural toxins.*, 4: 51- 52.
- Butler, T. (1902): "Notes on a feeding experiment to produce leukoencephalitis in a horse, with positive results" *Am. Vet. Rev.*, 26: 748- 751.
- Calnek, B.W.; Barnes, H.J.; Beard, C.W.; McDougald, L.R. and Saif, Y.M. (1997): "Diseases of poultry." 10th ed. Pp. 956- 957, Mosby- Wolfe publisher, USA.
- Caramelli, M.; Dondo, A.; Cortellazzi, G.C.; Visconti, A.; Minervini, F.; Doko, M. B. and Guarda, F. (1993): "Leukoencephalomalacia in the equine by funonisins: first report in Italy" *Ippologia*, 4(4): 49-56.
- Casteel, S.W.; Turk, J.R.; Cowart, R.P. and Rottinghaus, E.G. (1993): "Chronic toxicity of fumonisin in weanling pigs" *J. Vet. Diagn. Invest.*, 5 (3): 413- 417.
- Casteel, S. W.; Turk, J.R. and Rottinghaus, G. E. (1994): "Chronic effects of dietry fumonisin on the heart and pulmonary vasculature of swine" *Fundam. Appl. Toxicol.*, 23: 518-524.
- Chu, F.S. and Li, G.Y. (1994): "Simultaneous occurrence of fumonisin B₁ and other mycotoxins in mouldy corn collected from the people's Republic of China regions with high incidence of esophageal cancer." *Appl. Environ. Microbiol.*, 60: 847- 852.
- Colvin, B.M. and Harrison, L.R. (1992): "Fumonisin- induced pulmonary oedema and hydrothorax in swine" *Mycopathologia.*, 117: 79- 82.
- Colvin, B.M.; Cooley, A.J. and Beaver, R.W. (1993): "Fumonisin toxicosis in swine: Clinical and pathology findings." *J. Vet. Diagn. Invest.*, 5: 232- 241.
- Cumpecht, L.A.; Beasley, V.R.; Weigel, R.M.; Parker, H.M.; Tumbleson, M.E.; Bacon, C.W.; Meredith, F.I. and Haschek, W.M. (1998): "Development of fumonisin-induced hepatotoxicity and pulmonary oedema in orally dosed swine: Morphological and biochemical alterations" *Tocicologic. Pathology*, 26 (6): 777-788.
- Domench, J.; Boccas, B.; Pellegrin, F.; Laurent, D.; Kohler, F.; Magnol, J. and Lambert, C. (1985): "Equine leukoencephalomalacia in New Caledonia." *Vet. J.*, 62: 422.
- Edrington, T.S.; Kamps- Hotzapple, C.A.; Harvey, R.B.; Kubena, L.F.; Elissalde, M.H. and Rottinghaus, G.E. (1995):

- "Acute hepatic and renal toxicity in lambs dosed with fumonisin- containing culture material." *J. Anim. Sci.*, 73: 508-515.
- Espada, Y.; Ruiz de Gopegui, R.; Cuadradas, C. and Cabanes, F.J. (1994): "Fumoinis mycotoxicosis in broilers: Weights and serum chemistry modifications." *Avian Dis.*, 38: 454- 460.
- Espada, Y.; Ruiz de Gopegui, R.; Cuadradas, C. and Cabanes, F.J. (1997): "Fumoinis mycotoxicosis in broilers: Plasma proteins and coagulation modifications." *Avian Dis.*, 41: 73- 79.
- Fazekas, B. and Bajmicy, E. (1996): "Occurrence of equine Leukoencephalomalacia caused by fumonisin B₁ mycotoxin in Hungary." *Magyar Allatorvosok Lapja*, 51 (8): 484- 487.
- FDA, Food and Drug Administration (2000): "Background paper in support of fumonisin levels in animal feed." U.S. FDA center for veterinary medicine, June 6, 2000.
- FDA, Food and Drug Administration (2001a): "Background paper in support of fumonisin levels in animal feed: Excultive summary of this Scientific Support document." U.S. FDA center for veterinary medicine, November 9, 2001.
- FDA, Food and Drug Administration (2001b): "Background paper in support of fumonisin levels in corn and corn products intended for human consumption." U.S. FDA center for safety and applied nutrition, center for veterinary medicine, November 9, 2001.
- FDA, Food and Drug Administration (2001c): "Fumonisin levels in human foods and animal feeds." U.S. FDA center for food safety and applied nutrition center for veterinary medicine, November 9, 2001.
- Flynn, G.L. (1985): "Mechanism of percutaneous absorption from physicochemical evidence. In: Bronaugh, R.L. and Maibach, H.I. (eds) percutaneous absorption." New York, Marcel Dekker, PP. 17- 42.
- Gelderblom, W.C.; Jaskiewicz, K.; Marasas, W.F.; Thiel, P.G.; Horak, R.H. and Kriek, N.P. (1988): "Fumonisin- Novel mycotoxins with cancer- promoting activity produced by *Fusarium moniliforme*." *Appl. Environ. Microbiol.*, 54: 1806- 1811.
- Goel, S.; Lenz, S.D.; Lumlerdacha, S.; Lovell, R.T.; Shelby, R.A.; Li, M. and Kempanien, B.W. (1994): "Sphingolipid levels in catfish exposed to fumonisins." *The Toxicologist*, 14 (1): (Abstract 771).
- Graham, R. (1936): "Cornstalk disease investigations. Toxic encephalitis an non-virus encephalomyelitis of horses." *Vet. Med.*, 31: 46.
- Groves, S.D.; Zhang, L.; Chang, Y.S.; Ross, P.F.; Casper, H.; Norred, W.P.; You, W.C. and Fraumeeni, J.F.Jr. (1999): "Fusarium mycotoxins in corn and corn products in a high- risk area for gastric cancer in Shangdong Province." *China. J. Assoc. Off.Anal. Chem. International*, 82 (3): 657- 662.
- Gumprecht, L.A.; Marcucci, A.; Weigel, R.M.; Vesonder, R.F.; Riely, R.T.; Showker, J.L.; Beasley, V.R. and Haschek, W.M. (1995): "Effects of intravenous fumonisin B₁ in rabbits: nephrotoxicity and sphingolipis alterations." *Natural Toxins*, 3 (5): 395- 403.
- Gurung, N.K.; Rankins, D.L.Jr.; Shelby, R.A. and Goel, S. (1998): "Effects of fumonisin B₁-contaminated feeds on weanling Angora goats." *J. Anim. Sci.*, 76: 2863-2870.

- Hammer, P.; Bluthgen, A. and Walte, H.G. (1996): "Carry-over of fumonisin B₁ in the milk of lactating cows." *Milch-wissenschaft*, 51: 691- 695.
- Harrison, L.R.; Colvin, B.M.; Greene, J.T.; Newman, L.E. and Cole, J.R.Jr. (1990): "Pulmonary oedema and hydrothorax in swine produced by fumonisin B₁, a toxic metabolite of *Fusarium moniliforme*." *J. Vet. Diagn. Invest.*, 2: 217- 221.
- Haschek, W.M.; Motelinn, G.; Ness, D.K.; Harlin, K.S.; Hall, W.F.; Vesonder, R.; Peterson, R.E. and Beasley, V.R. (1992): "Characterization of fumonisin toxicity in orally and intravenously dosed swine." *Mycopathologia*, 117: 83- 96.
- Haschek, W.M.; Gumprecht, L.A.; Smith, G.W.; Parker, H.M.; Beasley, V.R. and Tumbleson, M.E. (1996): "Effects of fumonisins in swine." In: *Advances in swine biochemical research*. New York, Plenum Press, Pp. 99- 112.
- Haschek, W.M.; Motelin, G.; Ness, D.K.; Harlin, K.S.; Hall, W.F.; Vesonder, R.; Peterson, R.E. and Beasley, V.R. (1992): "Characterization of fumonisin toxicity in orally and intravenously dosed swine." *Mycopathologia*, 117: 83- 96.
- Henry, M.H.; Wyatt, R.D. and Fletcher, O.J. (2000): "The toxicity of purified fumonisin B₁ in broiler chicks." *Poult. Sci.*, 79 (10): 1378- 1384.
- Jaskiewicz, K.; Marasas, W.F. and Var de Walt, F.E. (1987): "Oesophageal and other main cancer patterns in four districts of Transkei, 1981- 1984. S. Afr." *Med. J.*, 72: 27- 30.
- Javed, T.; Bennet, G.A.; Richard, J.L.; Dombbrinkkurtzman, M.A.; Cote, L.M. and Buck, W.B. (1993): "Mortality in broiler chicks on feed attended with *Fusarium proliferatum* culture material or with purified fumonisin B₁ and moniliformin." *Mycopathologia*, 123: 171- 184.
- Jindal, N.; Mahipal, S.K. and Rottinghaus, G.E. (1999): "Occurrence of fumonisin B₁ in maize and poultry feeds in Haryana India." *Mycopathologia*, 148 (1): 37- 40.
- Jubb, K.V.F. and Kennedy, P.C. (1985): "Pathology of domestic animals." 2nd ed., New York, Academic Press.
- Kellerman, T.S.; Marasas, W.F.; Pienaar, J.G. and Nande, T.W. (1972): "A mycotoxicosis of equine caused by *Fusarium moniliforme* sheldon." *Onderst. J. Vet. Res.*, 39: 205- 208.
- Kellerman, T.S.; Marasas, W.F.; Thiel, P.G.; Gelderblom, W.C.; Cawood, M. and Coetzer, J.A. (1990): "Leukoencephalomalacia in two horses by oral dosing of fumonisin B₁." *Onderst. J. Vet. Res.*, 57: 269- 275.
- Klittich, C.J.; Laslie, J.F.; Nelson, P.E. and Marasas, W.F. (1997): "*Fusarium thapsinum* (*Gibberella thapsina*): A new species in section *Liseola* from sorghum." *Mycologia*, 89: 643- 652.
- Kobluk, C.N.; Ames, T.R. and Geor, R.J. (1995): "Horse, Diseases and Clinical Management." pp. 435-441, W.B. Saunders Company, Philadelphia, London, Toronto, Montreal, Sydney, Tokyo.
- Kriek, N.P.; Kellerman, T.S. and Marasas, W.F. (1981): "A comparative study of the toxicity of *Fusarium veticillioides* (= *F. moniliforme*) to horses, primates, pigs, sheep and rats." *Onderst. J. Vet. Res.*, 48: 129- 131.
- LaBorde, J.B.; Terry, K.K.; Howard, P.C.; Chen, J.J.; Collins, T.F.; Shackelford, M.E. and Hansen, D.K. (1997): "Lack of embryotoxicity of fumonisin B₁ in New-

- Zealand white rabbits." *Fundam. Appl. Toxicol.*, 40: 120- 128.
- Laurent, D.; Platzer, N.; Kohler, F.; Sauviat, M.P. and Pallegriin, F.(1989): "Macrofusine et micromoniline: Deux nouvelles mycotoxines isolees de mais infeste par *Fusarium moniliforme* sheld." *Microbiol. Alim. Nutr.*, 7: 9- 16.
- Ledoux, D.R.; Bermudez, A.J. and Rottinghaus, G.E. (1996): "Effects of feeding *Fusarium moniliforme* culture material, containing known levels of fumonisin B₁, in the young turkey poults." *Poult. Sci.*, 75 (12): 1472- 1478.
- Ledoux, D.R.; Brown, T.P., Weibking, T.S. and Rottinghaus, G.E. (1992): "Fumonisin toxicity in broiler chicks." *J. Vet. Diag. Invst.*, 4: 330- 333.
- Leeson, S.; Diaz, G.J. and Summers, J.D. (1995): "Poultry metabolic disorders and mycotoxins." PP. 299-309. University Books, Guelph, Ontario, Canada.
- Li, M.H.; Raverty, S.A. and Robinson, E.H. (1994): "Effects of dietary mycotoxins produced by the mold *Fusarium moniliforme* on channel catfish, *Ictalurus punctatus*." *J. of the World Agriculture Society*, 25 (4): 512- 516.
- Li, M.M. and Robinson, E.H. (1995): "Catfish growth not affected by low levels of mycotxins produced by the mold *Fusarium moniliforme*, new research indicates" *The Catfish Journal*, June, P.16.
- Logrieco, A.; Doko, M.B.; Moretti, A.; Frisullo, S. and Visconti, A. (1998): "Occurrence of fumonisin B₁ and B₂ in *Fusarium proliferatum* infected Asparagus plants." *J. Agric. Food Chem.*, 46: 5201- 5204.
- Lumlertdacha, S.; Lovell, R.T.; Shelby, R.A.; Lenz, S.D. and Kamppainen, B.W. (1995): "Growth hematology and histopathology of channel catfish, *Ictalurus punctatus*, fed toxins from *Fusarium moniliforme*." *Aquaculture*, 130 (2-3): 201- 218.
- Makaula, N.A.; Marasas, W.F.; Venter, F.S.; Badenhorst, C.J.; Bradshaw, D. and Swanevelder, S. (1996): "Oesophageal and other cancer patterns in four selected districts of Trankei, Southern Africa: 1985- 1990." *Afr. J. Health. Sci.*, 3:11-15.
- Marasas, W.F.; Kellerman, T.S.; Pienaar, J.G. and Naude, T.W. (1976): "Leukoencephalomalacia: A mycotoxicosis of Equidae caused by *Fusarium moniliforme* shaldon." *Ondest. J. Vet. Res.*, 43: 113- 122.
- Marasas, W.F.O.; Kellerman, T.S.; Gelderblom, W.C.A.; Coetzer, J.A.W.; Thiel, P.G. and Vander Lugh, J.J. (1988): "Leukoencephalomalacia in a horse induced by fumonisin B₁ isolated from *Fusarium moniliforme*." *Ondersteport J. Vet. Res.*, 55: 197- 203.
- Maragos, C.M. and Richard, J.L. (1994): "Quantitation and stability of fumonisins B₁ and B₂ in milk." *J. Assoc. Off.Anal. Chem. Int.*, 77: 1162- 1167.
- Masri, M.D.; Olcott, B.N.; Nicholson. S.S.; McClure, J.J.; Schimdt, S.P.; Freestone, J.F. and Kornagay, W.R. (1987): "Clinical, epidemiologic and pathologic evaluation of an outbreak of mycotoxin encephalomalacia in South Louisiana horses." *Proceedings of the American Association of Equine practitioners*, 367- 374.
- Mayes, P.A. (1988): "Metabolism of acylglycerols and sphingolipids. In: Murray, R.Y.; Ganner, D.K.; Mayes, P.A. and Rodwell, V.W. (eds.) *Harper's Biochemistry*. 21st "ed. Appleton and Lange Norwalk, Pp. 218- 225.

- McCue, P.M. (1989): "Equine leukoencephalomalacia." *Compend. Cotin. Educ. Pract. Vet.*, 11: 646- 651.
- Meister, U.; Symmank, H. and Dahlke, H. (1996): "Investigation and evaluation of the contamination of native and imported cereals with fumonisins. Z." *Lebensm. Unters. Forsch.*, 203: 528- 533.
- Merrill, A.H.; Schmelz, E.M.; Dillehay, D.L.; Spiegel, S.; Shayman, J.A.; Schroeder, J.J.; Riley, R.T.; Voss, K.A. and Want, E. (1997): "Sphingolipids- the enigmatic lipid class: biochemistry, physiology and pathophysiology." *Toxicol. Appl. Pharmacol.*, 142: 208- 225.
- Merrill, A.H.; Wang, E.; Gilehrst, D.G. and Riley, R.T. (1993): "Fumonins and other inhibitors of de novo sphingolipid biosynthesis." *Adv. Lipid Res.*, 26: 215- 234.
- Miller, J.D. (1999): "Factors affecting the occurrence of fumonisin in corn. Abstract of paper: International Conference on the Toxicology of Fumonisin." June 28- 30, 1999, Arlington, VA, P. 21.
- Mortelin, G.K.; Haschek, W.M.; Ness, D.K.; Hall, W.F.; Harlin, K.S.; Schaeffer, D.J. and Beasley, V.R. (1994): "Temporal and dose- response features in swine fed corn screenings contaminated with fumonisin mycotoxins." *Mycopathologia*, 126 (1): 27- 40.
- Musser, S.M. and Plattner, R.D. (1997): "Fumonisin composition in cultures of *Fusarium moniliforme*, *Fusarium proliferatum*, and *Fusarium nygami*." *J. Agric. Food Chem.*, 45: 1169- 1173.
- Norred, W.P.; Riely, R.T.; Meredith, F.J.; Bacon, C.W. and Voss, K.A. (1996): "Time- and dose- response effects of the mycotoxin, fumonasin B₁ on sphingoid base elevation in precision- cut rat liver and kidney slices." *Toxicol. In Vitro*, 10: 349- 358.
- Norred, W.P.; Voss, K.A.; Riely, R.T.; Meredith, F.I.; Bacon, C.W. and Merrill, A.H.Jr. (1998): "Mycotoxins and health hazards. Toxicological aspects and mechanism of action of fumonisins." *J. Toxicol. Sci.*, 23 (II) 160- 164.
- Norred, W.P.; Wang, E.; Hwangsoo, Y.; Riley, R.T. and Merrill, A.H.Jr. (1992): "In vitro toxicology of fumonisins and the mechanistic implications." *Mycopathologia*, 117: 73- 78.
- NTP, National Toxicology Program (1999): "Toxicology and carcinogenesis studies on fumonisin B₁ in F344/ N rats and B6CF1 mice. Technical Report Series, (496). NIH publication No. 99- 3955. National Institute of Health, Research Triangle Park, NC."
- Oswailer, C.D. (1999): "Fumonins. In: Howard, J.L. and Smith, R.A. (eds.) *Current Veterinary Therapy, Food Animal Practice*." Pp. 258- 259. W.B. Saunders, Co., Philadelphia, London, Toronto, Montreal, Sydney, Tokyo.
- Oswailer, G.D.; Kehri, M.E.; Stabel, J.R.; Thruston, J.R.; Ross, P.F. and Wilson, T.M. (1993): "Effects of fumonisin contaminated corn screenings on growth and health of feeder calves." *J. Anim. Sci.*, 71: 459- 466.
- Oswailer, G.D.; Ross, D.F.; Wilson, T.M.; Nelson, P.E.; Witte, S.T.; Carson, T.L.; Rice, L.G. and Nelson, H.A. (1992): "Characterization of an epizootic of pulmonary oedema in swine associated with fumonisin in corn screenings." *J. Vet. Diagn. Invest.*, 4: 53- 59.
- Placinta, C.M.; D'Mello, J.P. and McDonald, A.M. (1999): "A review of worldwide contamination of cereal grains and

- animal feed with fusarium mycotoxins." *Anim. Feed Sci. Technol.*, 78(1- 2): 21- 37.
- Plattner, R.D. (1995): "Detection of fumonisins produced in *Fusarium moniliforme* cultures by HPLC with electrospray MS and evaporative light scattering detectors." *Nat. Toxins*, 3: 294- 298.
- Plumlee, K.H. (1997): "Mycotoxicosis." In: Robenson, N.E. (ed.), *Current therapy in equine medicine*. P. 668, W.B. Saunders Co., USA.
- Plumlee, K.H. (1997): "Mycotoxins." In: N.E. Robeson (Ed.), *Current Therapy in Equine Medicine*, p. 668, W.B. Saunders Company, Philadelphia.
- Prelusky, D.B.; Trenholm, H.L. and Savard, M.E. (1994): "Pharmacokinetic fate of ¹⁴C- labelled fumonisin B₁ in swine." *Nat. Toxins*, 2: 73- 80.
- Prelusky, D.B.; Savard, M.E. and Trenholm, H.L. (1995): "Pilot study on the plasma pharmacokinetics of fumonisin B₁ in cows following a single dose by oral gavage or intravenous administration." *Nat. Toxins*, 3: 389- 394.
- Prelusky, D.B.; Miller, J.D. and Trenholm, H.L. (1996a): "Disposition of ¹⁴C- derived residues in tissues of pigs fed radiolabelled fumonisin B₁." *Food Addit. Contam.*, 13: 155- 162.
- Prelusky, D.B.; Trenholm, H.L.; Rotter, B.A.; Miller, J.D.; Savard, M.E.; Young, J.M. and Scott, P.M. (1996b): "Biological fate of fumonisin B₁ in food- producing animals." *Adv. Exp. Med. Biol.*, 392: 265- 278.
- Price, W.D.; Lovell, R.A.; McChesney, D.G. (1993): "Naturally occurring toxin in feedstuffs: Center for Veterinary Medicine Perspective." *J. Anm. Sci.*, 71: 2556- 2562.
- Qureshim M.A. and Hagler, W.M. (1992): "Effect of fumonisin B₁ exposure on chicken macrophage function in vitro." *Poult. Sci.*, 71: 104- 112.
- Rheeder, J.P.; Marasas, W.F.; Thiel, P.G.; Sydenham, E.W.; Shephard, G.S. and Van Schalkwyk, D.J. (1992): "*Fusarium moniliforme* and fumonisins in corn in relation to human esophageal cancer in Transkei." *Phytopathology*, 82: 353- 357.
- Rice, L.G. and Ross, P.F.(1994): "Methods for detection and quantitation of fumonisins in corn, cereal products and animal excreta." *J. Food Prot.*, 57: 536- 450.
- Richard, J.L.; Meerdink, G.; Maragos, C.M.; Tumbleson, M.; Bordson, G.; Rice, L.G. and Ross, P.F. (1996): "Absence of detectable fumonisins in the milk of cows fed *Fusarium proliferatum* Nirenberg culture material." *Mycopathologia*. 133: 123- 126.
- Riley, R.T.; An, N.H.; Showker, J.L.; Yoo, H.S.; Norred, W.P.; Chamberlain, W.J.; Wang, E.; Merrill, A.H.Jr.; Motelin, G.; Beasley, V.R. and Haschek, W.M. (1993): "Alteration of tissue and serum sphinganine to sphingosine ratio: An early biomarker of exposure to fumonisin- containing feeds in pigs." *Toxicol. Appl. Pharamacol.*, 118: 105- 112.
- Rooney, J.R. and Robertson, J.L. (1996): "Equine pathology." pp. 324- 325. Iowa state university press, Ames.
- Ross, P.F.; Ledet, A.E.; Owens, D.Z.; Ric, L.G.; Nelson, H.A.; Osweiler, G.D. and Wilson, T.M. (1993): "Experimental equine leukoencephalomalacia, toxic hepatitis and encephalopathy cause by corn naturally contaminated with fumonisine." *J. Vet. Diagn. Invest.* 5: 69-74.

- Ross, P.F.; Nelson, P.E.; Richard, J.L.; Osweiler, G.D.; Rice, L.G.; Plattner, R.D. and Wilson, T.M. (1990): "Production of fumonisins by *Fusarium moniliforme* and *Fusarium proliferatum* isolates associated with equine leukoencephalomalacia and a pulmonary edema syndrome in swine." *Appl. Environ. Microbiol.*, 56:3225- 3226.
- Ross, P.F.; Rice, L.G.; Plattner, R.D.; Osweiler, G.D.; Wilson, T.M.; Owens, D.L.; Nelson, H.A. and Richard, J.L. (1991): "Concentrations of fumonisin B₁ in feeds associated with animal health problems." *Mycopathologia.*, 114(3): 129- 135.
- Ross, P.F.; Rice, L.G.; Osweiler, G.D.; Nelson, P.E.; Richard, J.L. and Wilson, T.M. (1992): "A review and update of animal toxicosis associated with fumonisin-contaminated feeds and production of fumonisins by fusarium isolates." *Mycopathologia.*, 117: 109- 114.
- Ross, P.F.; Nelson, P.E.; Owens, D.L.; Rice, L.G.; Nelson, H.A. and Wilson, T.M. (1994): "Fumonisin B₂ in cultured *Fusarium proliferatum*, M- 6104, causes equine leukoencephalomalacia." *J. Vet. Diagn. Invest.*, 6: 263- 265.
- Rotter, B.A.; Thompson, B.K.; Prelusky, D.B.; Trenholm, H.L.; Stewart, B.; Miller, J.D. and Savard, M.E. (1996): "Response of growing swine to dietary exposure to pure fumonisin B₁ during an eight- week period: Growth and clinical parameters." *Nat. Toxins.*, 4 (1): 42- 50.
- Russell, M.A.; Eikenberry, J.G.; Scott, D.H. and Hop, W.D. (1993): "Moldy corn outbreak: the purdue CES Response. Proc. 13th Equine Nutr. and Physiol. Symp." Gainesville, Floride, pp. 286- 287.
- Schmidt, S.P. and Osborn, T.G. (1993): "Effects of endophyte- infected tall fescue on animal performance." *Agric. Ecosys. Environ.*, 44: 233- 262.
- Scott, P.M.; Delgado, T.; Prelusky, D.B.; Trenholm, H.L. and Miller, J.D. (1994): "Identification of fumonisin in milk." *J. Environ. Sci. Health*, B29: 989- 998.
- Scott, P.M.; Manhere, S.R.; Lawrence, G.A.; Daley, E.F. and Farber, J.M. (1995): "Fermentation of wort containing added ochratoxin A and Fumonisin B₁ and B₂". *Food Addit. Contam.*, 12: 31- 40.
- Shelby, R.A.; White, D.G. and Bauske, E.M. (1994): "Differential fumonisin production in maize hybrids." *Plant Dis.*, 78: 582- 584.
- Shephard, G.S.; Marasas, W.F.; Yazdanpanah, H.; Rahimian, H.; Safavi, N.; Zarghi, A.; Shafaati, A. and Rasekh, H.R. (2002): "Fumonisin B₁ in maize harvested in Iran during 1999." *Food Additives and Contaminants*, 19 (7): 676- 679.
- Shepherd, G.S.; Thiel, P.G.; Sydenham, E.W.; Alberts, J.F. and Cawood, M.E. (1994a): "Distribution and excretion of a single dose of the mycotoxin fumonisin B₁ in a non- human primate." *Food. Chem. Toxicol.*, 32: 735- 741.
- Shepherd, G.S.; Thiel, P.G.; Sydenham, E.W.; Vleggaar, R. and Alberts, J.F. (1994b): "Determination of the mycotoxin fumonisin B₁ and identification of its partially hydrolysed metabolites on the faeces on non- human." *Food Chem. Toxicol.*, 32: 23- 29.
- Shetty, P.H. and Bhat, R.V. (1997): "Natural occurrence of fumonisin B₁ and its Co-occurrence with aflatoxin B₁ in Indian sorghum, maize and poultry feeds." *J. Agric. Food Chem.*, 45: 2170- 2173.
- Smith, B.P. (2000): "Large animal internal medicine." 2nd ed. P. 1892. Mosby, Inc., USA.
- Smith, G.W.; Constable, P.D.; Tumbleson, M.E.; Rottinghaus, G.E. and Haschek,

- W.M. (1999): "Sequence of cardiovascular changes leading to pulmonary oedema in swine fed culture material containing fumonisin." *Am. J. Vet. Res.*, 60 (10): 1292- 1300.
- Smith, J.S. and Thakur, R.A. (1996): "Occurrence and fate of fumonisins in beef. In: L.S. Jackson, J.W. Devries and L.B. Bullerman (Ed.), *Fumonisin in food*." Pp. 39-55. Plenum Press, New York.
- Stack, J. (2003): "Fumonisin in corn." Nebraska Cooperative Extension NFO3-570 under plant diseases, C-17, field crop.
- Steyn, P.S. (1995): "Mycotoxins, general view, chemistry and structure." *Toxicol. Lett.*, 82- 83: 843- 851.
- Sydenham, E.W.; Shephard, G.S.; Thiel, P.G.; Marasas, W.F.O and Stockenstrom, S. (1991): "Fumonisin contamination of commercial corn-based human foodstuffs." *J. Agric. Food Chem.*, 39 (11): 2014-2018.
- Sydenham, E.W.; Marasas, W.F.; Shephard, G.S.; Thiel, P.G. and Hirooka, E.Y. (1992): "Fumonisin concentrations in Brazilian feeds associated with field outbreaks of confirmed and suspected animal mycotoxicosis." *J. Agric. Food Chem.*, 40 (6): 994-997.
- Tejada- Simon, M.V.; Marovatsanga, L.T. and Pastka, J.J. (1995): "Comparative detection of fumonisin by HPLC, ELISA and immunocytochemical localization in fusarium culture." *J. Food. Prot.*, 58: 666- 672.
- Thil, P.G.; Marasas, W.F.; Sydenham, E.W.; Shephard, G.S.; Gelderblom, W.C. and Nieuwenhuis, J.J. (1991a): "Survey of fumonisin production by fusarium species." *Appl. Environ. Microbiol.*, 57: 1089- 1093.
- Thiel, P.G.; Shephard, G.S.; Sydenham, E.W.; Marasas, W.F.O.; Nelson, P.E. and Wilson, T.M. (1991b): "Levels of fumonisin B₁ in feeds associated with confirmed cases of equine leukoencephalomalacia." *J. Agric. Food Chem.*, 39 (1): 109- 111.
- Thompson, F.N. and Stuedemann, J.A. (1993): "Pathophysiology of fescue toxicosis." *Agric. Ecosys. Environ.*, 44: 263- 281.
- Vincell, P. and Parker, G. (1999): "Mycotoxins in corn produced by *Fusarium* fungi." ID- 121- Electronic file created in 1999. Corn Growers Assoc., Kentucky, Frankfort.
- Vudathala, D.K.; Prelusky, D.B.; Ayroud, M.; Trenholm, H.L. and Miller, J.D. (1994): "Pharmacokinetic fate and pathological effects of ¹⁴C- fumonisin B₁ in laying hens." *Nat. Toxin.*, 2: 81- 88.
- Wang, E.; Norred, W.P.; Bacon, C.W.; Riley, R.T. and Merrill, A.H. (1991): "Inhibition of sphingolipid biosynthesis by fumonisins." *J. Biol. Chem.*, 266: 14486- 14490.
- Wang, E.; Ross, P.F.; Wilson, T.M.; Riley, R.T. and Merrill, A.H. (1992): "Increases in serum sphingosine and sphinganine and decreases in complex sphingolipids in poines given feed containing fumonisin, mycotoxins produced by *Fusarium moniliforme*." *J. Nutr.*, 122: 1706- 1716.
- Weibking, T.S.; Ledoux, D.R.; Bermudez, J.R. A.J.; Turk, G.E.; Rottinghaus, G.E.; Wang, E. and Merrill, A.H.Jr. (1993a): "Effects of feeding *Fusarium moniliforme* culture material containing known levels of fumonisin B₁, on the young broiler chick." *Poult. Sci.*, 72 (3): 456- 466.
- Weibking, T.S.; Ledoux, D.R.; Brown, D.R. and Rottinghaus, G.E. (1993b): "Fumonisin

- toxicity in turkey poults." *J. Vet. Diagn. Invest.*, 5: 75- 83.
- Weibking, T.; Ledoux, D.R.; Bermudez, A.J.; Turk, J.R. and Rottinghaus, G.E. (1995): "Effects on turkey poults of feeding *Fusarium moniliforme* M- 1325 culture material grown under different environmental conditions." *Avian. Dis.*, 39: 32- 38.
- Weidenborner, M. (2001): "Encyclopedia of food mycotoxins." Speinger publisher, Berlin, New York and London.
- WHO, World Health Organization (2000): "Fumonisin B₁. Environmental Health Criteria 219. International programme on chemical safety, WHO, Geneva."
- Wilkins, P.A.; Vaala, W.E.; Zivotofsky, D. and Twitchell, E.D. (1994): "A herd outbreak of equine leukoencephalomalacia." *Cornell. Veterinarian*, 84 (1): 53- 59.
- Wilson, B.J. and Maronport, R.R. (1971): "Causative fungus agent of leukoencephalomalacia in equine animals." *Vet. Rec.*, 88: 484- 486.
- Wilson, T.M.; Nelson, P.E.; Ryan, T.B.; Rouse, C.D.; Pittman, C.W.; Neal, T.P.; Porterfield, M.L. and Saunders, G.K. (1985): "Linking leukoencephalomalacia to commercial horse rations." *Vet. Med.*, 80: 63.
- Wilson, T.M.; Ross, P.F.; Owens, D.L.; Rice, L.G.; Green, S.A.; Jenkins, S.J. and Nelson, H.A. (1992): "Experimental reproduction of ELEM- A study to determine the minimum toxic dose in ponies." *Mycopathologia*, 117: 115- 120.
- Wilson, T.M.; Ross, P.F.; Rice, L.G.; Oswelner, G.D.; Nelson, H.A.; Owens, D.L.; Plattner, R.D.; Reggiardo, C.; Noon, J.H. and Pickrell, J.W. (1990): "Fumonisin B₁ levels associated with an epizootic of equine leukoencephalomalacia." *J. Vet. Diagnostic investigation*, 2 (3): 213- 216.

السموم الفطرية فى الأغذية والأعلاف

4- الفيومونيزينس

بدير إبراهيم عجاج

قسم بحوث الكيمياء والنقص الغذائى والسموم - معهد بحوث صحة الحيوان بالدقى - مركز البحوث الزراعية

السموم الفطرية هى مركبات كيميائية تنتجها أنواع من الفطريات التى تنمو على المواد الغذائية والأعلاف الحيوانية وتسبب أضرارا صحية لكل من الإنسان والحيوان. وتعتبر سموم الفيومونيزينس والتى تم اكتشافها حديثا ذات أهمية كبرى نظرا لإمكانية تواجدها بكميات كبيرة فى الأغذية والأعلاف (خاصة المحتوية على منتجات الذرة) الملوثة بفطريات الفيوزاريوم المفرزة لها.

ففى الإنسان لاحظ العلماء وجود علاقة بين إزدیاد معدل حدوث سرطان المرئىء وتواجد سموم الفيومونيزينس أو الفطريات المفرزة لها فى منتجات الذرة المستخدمة فى الغذاء فى بعض دول العالم. وفى الحيوان لوحظ كثرة حدوث حالات الأعراض العصبية والنفوق فى الخيول ، وحالات الأودیما الرؤیة والنفوق فى الخنازير فى العديد من دول العالم ومنها مصر. كما أثبت العلماء تجريبيا أن سموم الفيومونيزينس لها تأثيرات سمية أخرى فى العجول والأغنام والماعز والدجاج والبط والرومى والأسماك.