

Fuzzification of strongly and locally strongly compact spaces*

O. R. Sayed
Assiut University, Egypt
and
Adem Kiliçman
University Putra Malaysia, Malaysia
Received : June 2021. Accepted : July 2021

Abstract

In this paper, some characterizations of fuzzifying strong compactness are given, including characterizations in terms of nets and pre-subbases. Several characterizations of locally strong compactness in the framework of fuzzifying topology are introduced and the mapping theorems are obtained.

Keywords and Phrases: *Lukasiewicz logic; semantics; fuzzifying topology; fuzzifying compactness; strong compactness; fuzzifying locally compactness; locally strong compactness.*

2000 Mathematics Subject Classification: *54A40, 54B10, 54D30.*

*This paper was presented in the 6th International Conference for Young Scientists in Basic and Applied Sciences ICYS-BAS19, 27-30 March 2019, Hawaii Le Jardin Resort, Hurghada-Egypt.

1. Introduction and Preliminaries

In the last few years fuzzy topology, as an important research field in fuzzy set theory, has been developed into a quite mature discipline [8-10, 13-14, 29]. In contrast to classical topology, fuzzy topology is endowed with richer structure, to a certain extent, which is manifested with different ways to generalize certain classical concepts. So far, according to Ref. [9], the kind of topologies defined by Chang [4] and Goguen [6] is called the topologies of fuzzy subsets, and further is naturally called L -topological spaces if a lattice L of membership values has been chosen. Loosely speaking, a topology of fuzzy subsets (resp. an L -topological space) is a family τ of fuzzy subsets (resp. L -fuzzy subsets) of nonempty set X , and τ satisfies the basic conditions of classical topologies [12]. On the other hand, Höhle in [7] proposed the terminology L -fuzzy topology to be an L -valued mapping on the traditional powerset 2^X of X . The authors in [11, 13-14, 20] defined an L -fuzzy topology to be an L -valued mapping on the L -powerset L^X of X . In 1952, Rosser and Turquette [21] proposed emphatically the following problem: If there are many-valued theories beyond the level of predicates calculus, then what are the detail of such theories? As an attempt to give a partial answer to this problem in the case of point set topology, Ying in 1991-1993 [30-32] used a semantical method of continuous-valued logic to develop systematically fuzzifying topology. Briefly speaking, a fuzzifying topology on a set X assigns each crisp subset of X to a certain degree of being open, other than being definitely open or not. Roughly speaking, the semantical analysis approach transforms formal statements of interest, which are usually expressed as implication formulas in logical language, into some inequalities in the truth value set by truth valuation rules, and then these inequalities are demonstrated in an algebraic way and the semantic validity of conclusions is thus established. There are already more than 100 papers in fuzzifying topology published in the last two decades, I guess. But only a few papers can properly use the semantic method introduced in the original papers of Ying, which I strongly believe, can provide more delicate characterization of fuzzifying topological structure. So far, there has been significant research on fuzzifying topologies [1-3, 17-18, 23-24]. For example, Ying [33] introduced the concepts of compactness and established a generalization of Tychonoff's theorem in the framework of fuzzifying topology. In [24] the concept of local compactness in fuzzifying topology is introduced and some of its properties are established. Generalized open sets play a very important role in General Topology and they are

now the research topics of many topologists worldwide. Indeed a significant theme in General Topology and Real Analysis is the study of variously modified forms of continuity, separation axioms etc. by utilizing generalized open sets. One of the most well known notions and also an inspiration source is the notion of pre-open [15] sets introduced by Mashhour, Abd El-Monsef and El-Deeb in 1982. In 1984, the authors in [16] considered a strong versions of compact spaces and locally compact spaces defined in terms of pre-open subsets of a topological space. Alternative characterizations of such spaces were obtained. The concept of mixed fuzzy topological spaces is studied and investigated in some papers (see for example, [6, 21, 29, 32-35] In [2] the concepts of fuzzifying pre-open sets and fuzzifying pre-continuity were introduced and studied. Also, the authors in [3] introduced some concepts of fuzzifying pre-separation axioms and clarified the relations of these axioms with each other as well as the relations with other fuzzifying separation axioms. Furthermore, in [1], Abd El-Baki and Sayed characterized the concepts of fuzzifying pre-irresolute functions and used the finite intersection property to give a characterization of fuzzifying strong compact spaces. In this paper, the concepts of pre-base and pre-subbase of fuzzifying P -topology are introduced. Other characterizations of fuzzifying strong compactness are given, including characterizations in terms of nets and pre-subbase. Several characterizations of locally strong compactness in the framework of fuzzifying topology are introduced and the mapping theorems are obtained. Thus we fill a gap in the existing literature on fuzzifying topology. We use the terminologies and notations in [1-3, 30-33] without any explanation. We note that the set of truth values is the unit interval and we do often not distinguish the connectives and their truth value functions and state strictly our results on formalization as Ying does. We will use the symbol \otimes instead of the second "AND" operation \wedge as dot is hardly visible. This mean that

$$[P] \leq [\varphi \rightarrow \psi] \Leftrightarrow [P] \otimes [\varphi] \leq [\psi].$$

All of the contributions in General Topology in this paper which are not referenced may be original.

We procure some useful definitions and results which are useful in the rest of the present paper.

A unary fuzzy predicate $\tau_P : 2^X \rightarrow [0, 1]$, called fuzzifying pre-open [2], is given as follows:

$$A \in \tau_P := \forall x(x \in A \rightarrow x \in Int(Cl(A))) , \text{i.e., } \tau_P(A) = \bigwedge_{x \in A} Int(Cl(A))(x).$$

Similarly, A unary fuzzy predicate $\mathcal{F}_P : 2^X \rightarrow [0, 1]$, called fuzzifying pre-closed [2], is given as follows: $A \in \mathcal{F}_P := X - A \in \tau_P$. The fuzzifying pre-neighborhood system of a point $x \in X$ [2] is denoted by $N_x^{P^X}$ (or N_x^P) : $2^X \rightarrow [0, 1]$ and defined as $N_x^P(A) = \bigvee_{x \in B \subseteq A} \tau_P(B)$. The fuzzifying pre-closure of a set $A \subseteq X$ [2], denoted by $Cl_P \in [0, 1]^X$, is defined as $Cl_P(A)(x) = 1 - N_x^P(X - A)$. If (X, τ) is a fuzzifying topological space and $N(X)$ is the class of all nets in X , then the binary fuzzy predicates $\triangleright^P, \propto^P : N(X) \times X \rightarrow [0, 1]$ [23] are defined as

$$S \triangleright^P x := \forall A(A \in N_x^{P^X} \rightarrow S \tilde{\subset} A), \quad S \propto^P x := \forall A(A \in N_x^{P^X} \rightarrow S \bar{\subset} A),$$

where " $S \triangleright^P x$ ", " $S \propto^P x$ " stand for " S pre-converges to x ", " x is a pre-accumulation point of S ", respectively; and " $\tilde{\subset}$ ", " $\bar{\subset}$ " are the binary crisp predicates "almost in", "often in", respectively. The degree to which x is a pre-adherence point of S is $adh_P S(x) = [S \propto^P x]$. If (X, τ) and (Y, σ) are two fuzzifying topological spaces and $f \in Y^X$, the unary fuzzy predicates $C_P, I_P : Y^X \rightarrow [0, 1]$, called fuzzifying pre-continuity [2], fuzzifying pre-irresoluteness [1], are given as

$$C_P(f) := \forall B(B \in \sigma \rightarrow f^{-1}(B) \in \tau_P), \quad I_P(f) := \forall B(B \in \sigma_P \rightarrow f^{-1}(B) \in \tau_P),$$

respectively. Let Ω be the class of all fuzzifying topological spaces. A unary fuzzy predicate $T_2^P : \Omega \rightarrow [0, 1]$, called fuzzifying pre-Hausdorffness [3], is given as follows:

$$T_2^P(X, \tau) = \forall x \forall y((x \in X \wedge y \in X \wedge x \neq y) \rightarrow \exists B \exists C(B \in N_x^P \wedge C \in N_y^P \wedge B \cap C \equiv \emptyset)).$$

A unary fuzzy predicate $\Gamma : \Omega \rightarrow [0, 1]$, called fuzzifying compactness [33], is given as follows:

$$\Gamma(X, \tau) := (\forall \mathfrak{R})(K_o(\mathfrak{R}, X) \longrightarrow (\exists \varphi)((\varphi \leq \mathfrak{R}) \wedge K(\varphi, A) \otimes FF(\varphi)))$$

and if $A \subseteq X$, then $\Gamma(A) := \Gamma(A, \tau/A)$. For K, K_o (resp. \leq and FF) see [31, Definition 4.4] (resp. [31, Theorem 4.3] and [33, Definition 1.1 and Lemma 1.1]). A unary fuzzy predicate $fI : [0, 1]^{2^X} \rightarrow [0, 1]$, called fuzzy finite intersection property [33], is given as

$$fI(\mathfrak{R}) := \forall \varphi((\varphi \leq \mathfrak{R}) \wedge FF(\varphi) \rightarrow \exists x \forall B(B \in \varphi \rightarrow x \in B)).$$

A fuzzifying topological space (X, τ) is said to be fuzzifying P -topological space [1] if $\tau_P(A \cap B) \geq \tau_P(A) \wedge \tau_P(B)$. A binary fuzzy predicate

$K_P : [0, 1]^{2^X \times 2^X} \rightarrow [0, 1]$, called fuzzifying pre-open covering [1], is given as $K_P(\mathfrak{R}, A) := K(\mathfrak{R}, A) \otimes (\mathfrak{R} \subseteq \tau_P)$. A unary fuzzy predicate $\Gamma_P : \Omega \rightarrow [0, 1]$, called fuzzifying strongly compactness [1], is given as follows:

$$(X, \tau) \in \Gamma_P := (\forall \mathfrak{R})(K_P(\mathfrak{R}, X) \longrightarrow (\exists \varphi)((\varphi \leq \mathfrak{R}) \wedge K(\varphi, X) \otimes FF(\varphi)))$$

and if $A \subseteq X$, then $\Gamma_P(A) := \Gamma_P(A, \tau/A)$. It is obvious that

$$\Gamma_P(X, \tau) := \Gamma(X, \tau_P), \quad \Gamma_P(A, \tau/A) := \Gamma(A, \tau_P/A)$$

and

$$\models K_o(\mathfrak{R}, A) \longrightarrow K_P(\mathfrak{R}, A).$$

A fuzzifying strongly compact space is a generalization of strongly compact space [16]. A space X is said to be strongly compact if every pre-open cover X has a finite subcover.

A unary fuzzy predicate $LC : \Omega \rightarrow [0, 1]$, called fuzzifying locally compactness [24], is given as follows:

$$(X, \tau) \in LC := (\forall x)(\exists B)((x \in Int(B)) \otimes \Gamma(B, \tau/B)).$$

A space X is said to be locally strongly compact [16] if each point of X has a neighbourhood which is a strongly compact subspace.

2. Fuzzifying pre-base and pre-subbase

Definition 2.1. Let (X, τ) be a fuzzifying topological space and $\beta_P \subseteq \tau_P$. Then β_P is called a pre-base of τ_P if β_P fulfils the condition:

$$\models A \in N_x^{P^X} \rightarrow \exists B((B \in \beta_P) \wedge (x \in B \subseteq A)).$$

Example 2.1. Let $X = \{a, b, c\}$, and $I = [0, 1]$. Define a mapping $\tau : P(X) \rightarrow I$ on X as follows: $\tau(\emptyset) = \tau(X) = 1$, $\tau(\{a, c\}) = 0$, $\tau(\{a, b\}) = \frac{1}{5}$, $\tau(\{b, c\}) = \frac{1}{2}$, $\tau(\{a\}) = 0$, $\tau(\{b\}) = \frac{3}{4}$, $\tau(\{c\}) = \frac{1}{2}$. Then we can easily verify that τ is a fuzzifying topology. By calculating,

$$\tau_P(\emptyset) = \tau_P(X) = 1, \quad \tau_P(\{a, c\}) = \frac{1}{4}, \quad \tau_P(\{a, b\}) = 1, \quad \tau_P(\{b, c\}) = 1, \\ \tau_P(\{a\}) = \frac{1}{4}, \quad \tau_P(\{b\}) = \frac{3}{4}, \quad \tau_P(\{c\}) = 1. \quad \text{If we set } \beta_P = \tau_P, \text{ then} \\ N_x^P(A) = \bigvee_{x \in B \subseteq A} \tau_P(B) = \bigvee_{x \in B \subseteq A} \beta_P(B). \quad \text{Obviously, } \beta_P \text{ is a pre-base of}$$

τ_P by Definition 2.1.

Theorem 2.1. β_P is a pre-base of τ_P if and only if $\tau_P = \beta_P^{(\cup)}$, where

$$\beta_P^{(\cup)}(A) = \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda).$$

Proof. Suppose that β_P is a pre-base of τ_P . If

$$\bigcup_{\lambda \in \Lambda} B_\lambda = A,$$

then from Theorem 3.1 (1) (b) in [2],

$$\tau_P(A) = \tau_P\left(\bigcup_{\lambda \in \Lambda} B_\lambda\right) \geq \bigwedge_{\lambda \in \Lambda} \tau_P(B_\lambda) \geq \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda).$$

Consequently,

$$\tau_P(A) \geq \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda)$$

To prove that

$$\tau_P(A) \geq \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda)$$

we first prove

$$\tau_P(A) = \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \tau_P(B).$$

(Indeed, assume $\delta_x = \{B : x \in B \subseteq A\}$. Then, for any

$$f \in \prod_{x \in A} \delta_x, \bigcup_{x \in A} f(x) = A,$$

and further we have

$$\begin{aligned} \tau_P(A) &= \tau_P\left(\bigcup_{x \in A} f(x)\right) \\ &\geq \bigwedge_{x \in A} \tau_P(f(x)) \\ &\geq \bigvee_{\substack{f \in \prod_{x \in A} \delta_x \\ x \in A}} \bigwedge_{x \in A} \tau_P(f(x)) \\ &= \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \tau_P(B). \end{aligned}$$

Also,

$$\tau_P(A) \leq \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \tau_P(B).$$

Therefore,

$$\tau_P(A) = \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \tau_P(B).$$

Now, since

$$\begin{aligned} N_x^{PX}(A) &\leq \bigvee_{x \in B \subseteq A} \beta_P(B), \\ \tau_P(A) &= \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \tau_P(B) = \bigwedge_{x \in A} N_x^{PX}(A) \\ &\leq \bigwedge_{x \in A} \bigvee_{x \in B \subseteq A} \beta_P(B) = \bigvee_{f \in \prod_{x \in A} \delta_x} \bigwedge_{x \in A} \beta_P(f(x)). \end{aligned}$$

Then,

$$\tau_P(A) \leq \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda).$$

Therefore,

$$\tau_P(A) = \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda).$$

In the other side, we assume that

$$\tau_P(A) = \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = A} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda)$$

and we will show that β_P is a pre-base of τ_P , i.e., for any $A \subseteq X$, $N_x^{PX}(A) \leq \bigvee_{x \in B \subseteq A} \beta_P(B)$. Indeed, if $x \in B \subseteq A$ and $\bigcup_{\lambda \in \Lambda} B_\lambda = B$, then there exists $\lambda_0 \in \Lambda$ such that $x \in B_{\lambda_0}$ and

$$\bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda) \leq \beta_P(B_{\lambda_0}) \leq \bigvee_{x \in B \subseteq A} \beta_P(B).$$

Therefore,

$$\begin{aligned} N_x^{PX}(A) &= \bigvee_{x \in B \subseteq A} \tau_P(B) \\ &= \bigvee_{x \in B \subseteq A} \bigvee_{\bigcup_{\lambda \in \Lambda} B_\lambda = B} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda) \end{aligned}$$

$$\leq \bigvee_{x \in B \subseteq A} \beta_P(B).$$

□

Theorem 2.2. Let $\beta_P : 2^X \rightarrow [0, 1]$. Then β_P is a pre-base for some fuzzifying P -topology τ_P if and only if it has the following properties:

- (1) $\beta_P^{(\cup)}(X) = 1$;
- (2) $\models (A \in \beta_P) \wedge (B \in \beta_P) \wedge (x \in A \cap B) \rightarrow \exists C((C \in \beta_P) \wedge (x \in C \subseteq A \cap B))$.

Proof. If β_P is a pre-base for some fuzzifying P -topology τ_P , then $\tau_P(X) = \beta_P^{(\cup)}(X)$. Clearly, $\beta_P^{(\cup)}(X) = 1$. In addition, if $x \in A \cap B$, then

$$\begin{aligned} \beta_P(A) \wedge \beta_P(B) \leq \tau_P(A) \wedge \tau_P(B) &\leq \tau_P(A \cap B) \leq N_x^{P_X}(A \cap B) \\ &\leq \bigvee_{x \in C \subseteq A \cap B} \beta_P(C). \end{aligned}$$

Conversely, if β_P satisfies (1) and (2), then we have τ_P is a fuzzifying P -topology. In fact, $\tau_P(X) = 1$. For any $\{A_\lambda : \lambda \in \Lambda\} \subseteq P(X)$, we set

$$\delta_\lambda = \left\{ \{B_{\Phi_\lambda} : \Phi_\lambda \in \Lambda_\lambda\} : \bigcup_{\Phi_\lambda \in \Lambda_\lambda} B_{\Phi_\lambda} = A_\lambda \right\}.$$

Then, for any

$$f \in \prod_{\lambda \in \Lambda} \delta_\lambda, \quad \bigcup_{\lambda \in \Lambda} \bigcup_{B_{\Phi_\lambda} \in f(\lambda)} B_{\Phi_\lambda} = \bigcup_{\lambda \in \Lambda} A_\lambda.$$

Therefore,

$$\begin{aligned} \tau_P \left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) &= \bigvee_{\substack{\Phi \in \Lambda \\ B_\Phi = \bigcup_{\lambda \in \Lambda} A_\lambda}} \bigwedge_{\Phi \in \Lambda} \beta_P(B_\Phi) \\ &\geq \bigvee_{\substack{f \in \prod_{\lambda \in \Lambda} \delta_\lambda \\ \lambda \in \Lambda}} \bigwedge_{\lambda \in \Lambda} \bigwedge_{B_{\Phi_\lambda} \in f(\lambda)} \beta_P(B_{\Phi_\lambda}) \\ &\geq \bigwedge_{\lambda \in \Lambda} \bigvee_{\{B_{\Phi_\lambda} : \Phi_\lambda \in \Lambda_\lambda\} \in \delta_\lambda} \bigwedge_{\Phi_\lambda \in \Lambda_\lambda} \beta_P(B_{\Phi_\lambda}) \\ &= \bigwedge_{\lambda \in \Lambda} \tau_P(A_\lambda). \end{aligned}$$

Finally, we need to prove that

$$\tau_P(A \cap B) \geq \tau_P(A) \wedge \tau_P(B).$$

If $\tau_P(A) > t, \tau_P(B) > t$, then there exists

$$\{B_{\lambda_1} : \lambda_1 \in \Lambda_1\}, \{B_{\lambda_2} : \lambda_2 \in \Lambda_2\}$$

such that $\bigcup_{\lambda_1 \in \Lambda_1} B_{\lambda_1} = A, \bigcup_{\lambda_2 \in \Lambda_2} B_{\lambda_2} = B$

and for any $\lambda_1 \in \Lambda_1, \beta_P(B_{\lambda_1}) > t$,

for any $\lambda_2 \in \Lambda_2, \beta_P(B_{\lambda_2}) > t$. Now, for any $x \in A \cap B$, there exists $\lambda_{1x} \in \Lambda_1, \lambda_{2x} \in \Lambda_2$ such that $x \in B_{\lambda_{1x}} \cap B_{\lambda_{2x}}$. From the assumption, we know that

$$t < \beta_P(B_{\lambda_{1x}}) \wedge \beta_P(B_{\lambda_{2x}}) \leq \bigvee_{x \in C \subseteq B_{\lambda_{1x}} \cap B_{\lambda_{2x}}} \beta_P(C),$$

and furthermore, there exists C_x such that

$$x \in C_x \subseteq B_{\lambda_{1x}} \cap B_{\lambda_{2x}} \subseteq A \cap B, \beta_P(C_x) > t.$$

Since

$$\bigcup_{x \in A \cap B} C_x = A \cap B,$$

we have

$$t \leq \bigwedge_{x \in A \cap B} \beta_P(C_x) \leq \bigvee_{\substack{\bigcup_{\lambda \in \Lambda} B_\lambda = A \cap B \\ \lambda \in \Lambda}} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda) = \tau_P(A \cap B).$$

Now, let $\tau_P(A) \wedge \tau_P(B) = k$. For any natural number n , we have

$$\tau_P(A) > k - \frac{1}{n}, \tau_P(B) > k - \frac{1}{n}$$

and so $\tau_P(A \cap B) \geq k - \frac{1}{n}$. Therefore $\tau_P(A \cap B) \geq k = \tau_P(A) \wedge \tau_P(B)$. \square

Definition 2.2. $\varphi_P : 2^X \rightarrow [0, 1]$ is called a pre-subbase of τ_P if $\varphi_P^{(\cap)}$ is a pre-base of τ_P , where

$$\varphi_P^{(\cap)}\left(\bigcap_{\lambda \in \Lambda} B_\lambda\right) = \bigvee_{\substack{\bigcap_{\lambda \in \Lambda} B_\lambda = A \\ \lambda \in \Lambda}} \bigwedge_{\lambda \in \Lambda} \varphi_P(B_\lambda),$$

$\{B_\lambda : \lambda \in \Lambda\} \sqsubset P(X)$ with " \sqsubset " standing for "a finite subset of".

Theorem 2.3. $\varphi_P : 2^X \rightarrow [0, 1]$ is a pre-subbase of some fuzzifying P -topology if and only if $\varphi_P^{(\cup)}(X) = 1$.

Proof. We only demonstrate that $\varphi_P^{(\cap)}$ satisfies the second condition of Theorem 2.2, and others are obvious. In fact

$$\begin{aligned}
& \varphi_P^{(\cap)}(A) \wedge \varphi_P^{(\cap)}(B) \\
&= \left(\bigcap_{\lambda_1 \in \Lambda_1} \bigvee_{B_{\lambda_1}=A} \bigwedge_{\lambda_1 \in \Lambda_1} \varphi_P(B_{\lambda_1}) \right) \wedge \left(\bigcap_{\lambda_2 \in \Lambda_2} \bigvee_{B_{\lambda_2}=B} \bigwedge_{\lambda_2 \in \Lambda_2} \varphi_P(B_{\lambda_2}) \right) \\
&= \bigvee_{\substack{\lambda_1 \in \Lambda_1 \\ B_{\lambda_1}=A}} \bigvee_{\substack{\lambda_2 \in \Lambda_2 \\ B_{\lambda_2}=B}} \left(\bigwedge_{\lambda_1 \in \Lambda_1} \varphi_P(B_{\lambda_1}) \right) \wedge \left(\bigwedge_{\lambda_2 \in \Lambda_2} \varphi_P(B_{\lambda_2}) \right) \\
&\leq \bigvee_{\substack{\lambda \in \Lambda \\ B_\lambda=A \cap B}} \left(\bigwedge_{\lambda \in \Lambda} \varphi_P(B_\lambda) \right) \\
&= \varphi_P^{(\cap)}(A \cap B).
\end{aligned}$$

Therefore, if $x \in A \cap B$, then

$$\varphi_P^{(\cap)}(A) \wedge \varphi_P^{(\cap)}(B) \leq \varphi_P^{(\cap)}(A \cap B) \leq \bigvee_{x \in C \subseteq A \cap B} \varphi_P^{(\cap)}(C).$$

□

3. Fuzzifying strong compact spaces

Theorem 3.1. Let (X, τ) be a fuzzifying topological space, φ_P be a pre-subbase of τ_P , and

$$(\beta_1) := (\forall \mathfrak{R})(K_{\varphi_P}(\mathfrak{R}, X) \rightarrow \exists \varphi((\varphi \leq \mathfrak{R}) \wedge K(\varphi, X) \otimes FF(\varphi))), \text{ where } K_{\varphi_P}(\mathfrak{R}, X) := K(\mathfrak{R}, X) \otimes (\mathfrak{R} \subseteq \varphi_P);$$

$$(\beta_2) := (\forall S)((S \text{ is a universal net in } X) \rightarrow \exists x((x \in X) \wedge (S \triangleright^P x)));$$

$$(\beta_3) := (\forall S)((S \in N(X) \rightarrow (\exists T)(\exists x)((T < S) \wedge (x \in X) \wedge (T \triangleright^P x))),$$

where " $T < S$ " stands for " T is a subnet of S ";

$$(\beta_4) := (\forall S)((S \in N(X) \rightarrow \neg(adh_P S \equiv \phi));$$

$$(\beta_5) := (\forall \mathfrak{R})(\mathfrak{R} \in [0, 1]^{2^X} \wedge \mathfrak{R} \subseteq \mathcal{F}_P \otimes fI(\mathfrak{R}) \rightarrow \exists x \forall A(A \in \mathfrak{R} \rightarrow x \in A)).$$

Then $\models (X, \tau) \in \Gamma_P \leftrightarrow \beta_i \ i = 1, 2, \dots, 5.$

Proof. (1) Since $\varphi_P \subseteq \tau_P$, $[\mathfrak{R} \subseteq \varphi_P] \leq [\mathfrak{R} \subseteq \tau_P]$ for any $\mathfrak{R} : 2^X \rightarrow [0, 1]$, then we have $[K_{\varphi_P}(\mathfrak{R}, X)] \leq [K_P(\mathfrak{R}, X)]$. Therefore $\Gamma_P(X, \tau) \leq [\beta_1]$.

(2) $[\beta_2] = \bigwedge \{ \bigvee_{x \in X} [S \triangleright^P x] : (S \text{ is a universal net in } X) \}$.

(2.1) Assume X is finite. We set $X = \{x_1, \dots, x_m\}$. For any universal net S in X , there exists $i_o \in \{1, \dots, m\}$ with $S \tilde{\subset} \{x_{i_o}\}$. In fact, if not, then for any $i \in \{1, \dots, m\}$, $S \tilde{\not\subset} \{x_{i_o}\}$, $S \tilde{\subset} X - \{x_{i_o}\}$ and $S \tilde{\subset} \bigcap_{i=1}^m (X - \{x_i\}) = \phi$, a contradiction. Therefore $x_{i_o} \notin A$ and $N_{x_{i_o}}^P(A) = 0$ (see[2],Theorem 4.2 (1)) provided $S \tilde{\not\subset} A$, and furthermore $[S \triangleright^P x_{i_o}] = \bigwedge_{S \tilde{\not\subset} A} (1 - N_{x_{i_o}}^P(A)) = 1$.

Therefore $[\beta_2] = 1 \geq [\beta_1]$.

(2.2) In general, to prove that $[\beta_1] \leq [\beta_2]$ we prove that for any $\lambda \in [0, 1]$, if $[\beta_2] < \lambda$, then $[\beta_1] < \lambda$. Assume for any $\lambda \in [0, 1]$, $[\beta_2] < \lambda$. Then, there exists a universal net S in X such that $\bigvee_{x \in X} [S \triangleright^P x] < \lambda$ and for

any $x \in X$, $[S \triangleright^P x] = \bigwedge_{S \tilde{\not\subset} A} (1 - N_x^P(A)) < \lambda$, i.e., there exists $A \subseteq X$

with $S \tilde{\not\subset} A$ and $N_x^P(A) > 1 - \lambda$. Since φ_P is a pre-subbase of τ_P , $\varphi_P^{(\cap)}$ is a pre-base of τ_P and from Definition 2.1, we have $\bigvee_{x \in B \subseteq A} \varphi_P^{(\cap)}(B) \geq$

$N_x^P(A) > 1 - \lambda$, i.e., there exists $B \subseteq A$ such that $x \in B \subseteq A$ and

$\bigvee \left\{ \min_{\lambda \in \Lambda} \varphi_P(B_\lambda) : \bigcap_{\lambda \in \Lambda} B_\lambda = B, B_\lambda \subseteq X, \lambda \in \Lambda \right\} = \varphi_P^{(\cap)}(B) > 1 - \lambda$, where

Λ is finite. Therefore, there exists a finite set Λ and $B_\lambda \subseteq X (\lambda \in \Lambda)$ such that $\bigcap_{\lambda \in \Lambda} B_\lambda = B$ and for any $\lambda \in \Lambda$, $\varphi_P(B_\lambda) > 1 - \lambda$. Since $S \tilde{\not\subset} A$ and Λ is

finite, there exists $\lambda(x) \in \Lambda$ such that $S \tilde{\not\subset} B_{\lambda(x)}$. We set

$$\mathfrak{R}_o(B_{\lambda(x)}) = \bigvee_{x \in X} \varphi_P(B_{\lambda(x)}).$$

If $\wp \leq \mathfrak{R}_o$, then for any $\delta > 0$, $\wp_\delta \subseteq \{B_{\lambda(x)} : x \in X\}$. Consequently, for any $B \in \wp_\delta$, $S \tilde{\not\subset} B$ and $S \tilde{\subset} B^c$ because S is a universal net. If

$$[FF(\wp)] = 1 - \inf \{ \delta \in [0, 1] : F(\wp_\delta) \} = t,$$

then for any $n \in w$ (the non-negative integer),

$$\inf \{ \delta \in [0, 1] : F(\wp_\delta) \} < 1 - t + \frac{1}{n},$$

and there exists $\delta_o < 1 - t + \frac{1}{n}$ such that $F(\wp_{\delta_o})$. If $\delta_o = 0$, then $P(X) = \wp_{\delta_o}$ is finite and it is proved in (2.1). If $\delta_o > 0$, then for any $B \in \wp_{\delta_o}$, $S \tilde{\subset} B^c$.

Since $F(\wp_{\delta_0})$, we have

$$S\tilde{\subset} \bigcap \{B^c : B \in \wp_{\delta_0}\} \neq \phi.$$

i.e., $\bigcup \wp_{\delta_0} \neq X$ and there exist $x_0 \in X$ such that for any $B \in \wp_{\delta_0}, x_0 \notin B$. Therefore, if $x_0 \in B$, then $B \notin \wp_{\delta_0}$, i.e., $\wp(B) < \delta_0$,

$$K(\wp, X) = \bigwedge_{x \in X} \bigvee_{x \in B} \wp(B) \leq \bigvee_{x_0 \in B} \wp(B) \leq \delta_0 < 1 - t + \frac{1}{n}.$$

Let $n \rightarrow \infty$. We obtain $K(\wp, X) \leq 1 - t$ and $[K(\wp, X) \otimes FF(\wp)] = 0$. In addition, $[K_{\varphi_P}(\mathfrak{R}_0, X)] \geq 1 - \lambda$. In fact, $[\mathfrak{R}_0 \subseteq \varphi_P] = 1$ and

$$K(\mathfrak{R}_0, X) = \bigwedge_{x \in X} \bigvee_{x \in B} \mathfrak{R}_0(B) \geq \bigwedge_{x \in X} \mathfrak{R}_0(B_{\lambda(x)}) \geq \bigwedge_{x \in X} \varphi_P(B_{\lambda(x)}) \geq 1 - \lambda$$

because $x \in B_{\lambda(x)}$. Now, we have

$$\begin{aligned} [\beta_1] &= [(\forall \mathfrak{R})(K_{\varphi_P}(\mathfrak{R}, X) \rightarrow \exists \wp((\wp \leq \mathfrak{R}) \wedge K(\wp, X) \otimes FF(\wp)))] \\ &\leq [K_{\varphi_P}(\mathfrak{R}_0, X) \rightarrow \exists \wp((\wp \leq \mathfrak{R}_0) \wedge K(\wp, X) \otimes FF(\wp))] \\ &= \min(1, 1 - K_{\varphi_P}(\mathfrak{R}_0, X) + \bigvee_{\wp \leq \mathfrak{R}_0} [K(\wp, X) \otimes FF(\wp)]) \leq \lambda. \end{aligned}$$

Since λ is arbitrary, we have $[\beta_1] \leq [\beta_2]$.

(3) It is immediate that $[\beta_2] \leq [\beta_3]$.

(4) To prove that $[\beta_3] \leq [\beta_4]$, first we prove that

$$[\exists T ((T < S) \wedge (T \triangleright^P x))] \leq [S \propto^P x],$$

where

$$[\exists T ((T < S) \wedge (T \triangleright^P x))] = \bigvee_{T < S} \bigwedge_{T \tilde{\zeta} A} (1 - N_x^P(A))$$

and

$$[S \propto^P x] = \bigwedge_{S \tilde{\zeta} A} (1 - N_x^P(A)).$$

Indeed, for any $T < S$ one can deduce $\{A : S \tilde{\zeta} A\} \subseteq \{A : T \tilde{\zeta} A\}$ as follows. Suppose $T = S \circ K$. If $S \tilde{\zeta} A$, then there exists $m \in D$ such that $S(m) \notin A$ when $n \geq m$, where \geq directs the domain D of S . Now, we will show that $T \tilde{\zeta} A$. If not, then there exists $p \in E$ such that $T(q) \in A$ when $q \geq p$, where \geq directs the domain E of T . Moreover, there exists $n_1 \in E$ such that

$K(n_1) \geq m$ because $T < S$, and there exists $n_2 \in E$ such that $n_2 \geq n_1, p$ because (E, \geq) is directed. So, $K(n_2) \geq K(n_1) \geq m$, $S \circ K(n_2) \notin A$ and $S \circ K(n_2) = T(n_2) \in A$. They are contrary. Hence,

$$\{A : S\tilde{\nabla}A\} \subseteq \{A : T\tilde{\nabla}A\}.$$

Therefore,

$$\begin{aligned} [\exists T ((T < S) \wedge (T \triangleright^P x))] &= \bigvee_{T < S} \bigwedge_{T\tilde{\nabla}A} (1 - N_x^P(A)) \\ &= \bigvee_{T < S} \bigwedge_{\{A : T\tilde{\nabla}A\}} (1 - N_x^P(A)) \\ &\leq \bigwedge_{\{A : S\tilde{\nabla}A\}} (1 - N_x^P(A)) \\ &= \bigwedge_{S\tilde{\nabla}A} (1 - N_x^P(A)) = [S \propto^P x]. \end{aligned}$$

Therefore, for any $x \in X$ and $S \in N(X)$ we have

$$\begin{aligned} [\beta_3] &= \bigwedge_{S \in N(X)} \bigvee_{x \in X} [\exists T ((T < S) \wedge (T \triangleright^P x))] \\ &\leq \bigwedge_{S \in N(X)} \bigvee_{x \in X} [S \propto^P x] = \bigwedge_{S \in N(X)} \neg \left(\bigwedge_{x \in X} (1 - [S \propto^P x]) \right) \\ &= \bigwedge_{S \in N(X)} [\neg(\text{adh}_P S \equiv \phi)] = [\beta_4]. \end{aligned}$$

(5) We want to show that $[\beta_4] \leq [\beta_5]$. For any $\mathfrak{R} : 2^X \rightarrow [0, 1]$, assume $[fI(\mathfrak{R})] = \lambda$. Then for any $\delta > 1 - \lambda$, if $A_1, \dots, A_n \in \mathfrak{R}_\delta$, $A_1 \cap A_2 \cap \dots \cap A_n \neq \phi$. In fact, we set $\wp(A_i) = \bigvee_{i=1}^n \mathfrak{R}(A_i)$. Then $\wp \leq \mathfrak{R}$ and $FF(\wp) = 1$. By putting $\varepsilon = \lambda + \delta - 1 > 0$, we obtain

$$\lambda - \varepsilon < \lambda \leq [FF(\wp) \rightarrow (\exists x)(\forall B)(B \in \wp \rightarrow x \in B)] = \bigvee_{x \in X} \bigwedge_{x \notin B} (1 - \wp(B)).$$

There exists $x_o \in X$ such that $\lambda - \varepsilon < \bigwedge_{x_o \notin B} (1 - \wp(B))$, $x_o \notin B$ implies $\wp(B) < 1 - \lambda + \varepsilon = \delta$ and $x_o \in \bigcap \wp_\delta = A_1 \cap A_2 \cap \dots \cap A_n$. Now, we set $\wp_\delta = \{A_1 \cap A_2 \cap \dots \cap A_n : n \in N, A_1, \dots, A_n \in \mathfrak{R}_\delta\}$ and $S : \wp_\delta \rightarrow X, B \mapsto$

$x_B \in B, B \in \vartheta_\delta$ and know that $(\vartheta_\delta, \subseteq)$ is a directed set and S is a net in X . Therefore,

$$[\beta_4] \leq [\neg(adh_P S \equiv \phi)] = \bigvee_{x \in X} \bigwedge_{S \tilde{\zeta} A} (1 - N_x^P(A)).$$

Assume that $[\mathfrak{R} \subseteq F_P] = \mu$. Then, for any $B \in P(X)$, $\mathfrak{R}(B) \leq 1 + F_P(B) - \mu$, and

$$\begin{aligned} [\mathfrak{R} \subseteq F_P \otimes fI(A) \rightarrow (\exists x)(\forall A)((A \in \mathfrak{R}) \rightarrow x \in A)] \\ = \min(1, 2 - \mu - \lambda + \bigvee_{x \in X} \bigwedge_{x \notin A} (1 - \mathfrak{R}(A))). \end{aligned}$$

Therefore, it suffices to show that for any $x \in X$,

$$\bigwedge_{S \tilde{\zeta} A} (1 - N_x^P(A)) \leq 2 - \mu - \lambda + \bigwedge_{x \notin A} (1 - \mathfrak{R}(A)),$$

i.e.,

$$\bigvee_{x \notin A} \mathfrak{R}(A) \leq 2 - \mu - \lambda + \bigvee_{S \tilde{\zeta} A} N_x^P(A)$$

for some $\delta > 1 - \lambda$. For any $t \in [0, 1]$, if $\bigvee_{x \notin A} \mathfrak{R}(A) > t$, then there exists A_o such that $x_o \notin A_o$ and $\mathfrak{R}(A_o) > t$.

Case 1. $t \leq 1 - \lambda$, then $t \leq 2 - \mu - \lambda + \bigvee_{S \tilde{\zeta} A} N_x^P(A)$.

Case 2. $t > 1 - \lambda$. Here we set $\delta = \frac{1}{2}(t + 1 - \lambda)$ and have $A_o \in \mathfrak{R}_\delta, A_o \in \vartheta_\delta$. In addition,

$$t < \mathfrak{R}(A_o) \leq 1 + F_P(A_o) - \mu, \quad t + \mu - 1 \leq F_P(A_o) = \tau_P(A_o^c).$$

Since $A_o \in \vartheta_\delta$, we know that $S_B \in A_o$, i.e., $S_B \notin A_o^c$ when $B \subseteq A_o$ and $S \tilde{\zeta} A_o^c$. Therefore,

$$2 - \mu - \lambda + \bigvee_{S \tilde{\zeta} A} N_x^P(A) \geq 2 - \mu - \lambda + N_x^P(A_o^c) \geq 2 - \mu - \lambda + \tau_P(A_o^c) \geq t + (1 - \lambda) \geq t.$$

By noticing that t is arbitrary, we have completed the proof.

(6) To prove that $[\beta_5] = [(X, \tau) \in \Gamma_P]$ [1, Theorem 6]. □

The above theorem is a generalization of the following corollary.

Corollary 3.1. *The following are equivalent for a topological space (X, τ) .*

- (a) X is a strong compact space.
- (b) Every cover of X by members of a pre-subbase of τ_P has a finite sub-cover.
- (c) Every universal net in X pre-converges to a point in X .
- (d) Each net in X has a subnet that pre-converges to some point in X .
- (e) Each net in X has a pre-adherent point.
- (f) Each family of pre-closed sets in X that has the finite intersection property has a non-void intersection.

The equivalence (a) , (b) and (c) were given in [16].

Definition 3.1. *Let $\{(X_s, \tau_s) : s \in S\}$ be a family of fuzzifying topological spaces, $\prod_{s \in S} X_s$ be the cartesian product of $\{X_s : s \in S\}$ and $\varphi = \{p_s^{-1}(U_s) : s \in S, U_s \in P(X_s)\}$, where $p_t : \prod_{s \in S} X_s \rightarrow X_t (t \in S)$ is a projection. For $\Phi \subseteq \varphi$, $S(\Phi)$ stands for the set of indices of elements in Φ . The pre-base $\beta_P : \prod_{s \in S} X_s \rightarrow [0, 1]$ of $\prod_{s \in S} (\tau_P)_s$ is defined as*

$$V \in \beta_P := (\exists \Phi)(\Phi \sqsubset \varphi \wedge (\bigcap_{s \in S(\Phi)} \Phi = V)) \rightarrow \forall s(s \in S(\Phi) \rightarrow V_s \in (\tau_P)_s),$$

i.e.,

$$\beta_P(V) = \bigvee_{\Phi \sqsubset \varphi, \bigcap_{s \in S(\Phi)} \Phi = V} \bigwedge_{s \in S(\Phi)} (\tau_P)_s(V_s).$$

Example 3.1. *Let (X, τ) and τ_P be just as in Example 2.1. Define a mapping $\varsigma : P(Y) \rightarrow I$ on Y as follows: $\varsigma(\emptyset) = \varsigma(Y) = 1$, where $Y = \{d\}$, then ς is a fuzzifying topology and $\varsigma_P(\emptyset) = \varsigma_P(Y) = 1$. Hence, $Y \times X = \{(d, a), (d, b), (d, c)\}$, so $\varphi = \{\emptyset, X \times Y, \{(d, a)\}, \{(d, b)\}, \{(d, c)\}, \{(d, a), (d, b)\}, \{(d, b), (d, c)\}, \{(d, a), (d, c)\}\}$. By calculating, $\beta_P(\emptyset) = 1, \beta_P(X \times Y) = 1, \beta_P(\{(d, a)\}) = \frac{1}{4}, \beta_P(\{(d, b)\}) = \frac{3}{4}, \beta_P(\{(d, c)\}) = \frac{1}{2}, \beta_P(\{(d, a), (d, c)\}) = \frac{1}{4}, \beta_P(\{(d, a), (d, b)\}) = \frac{1}{2}, \beta_P(\{(d, b), (d, c)\}) = 1$. According to Theorem 2.1, we can easily obtain $\beta_P^{(U)} = \beta_P$, so $\tau_P \times \varsigma_P = \beta_P$.*

Definition 3.2. *Let $(X, \tau), (Y, \sigma)$ be two fuzzifying topological space. A unary fuzzy predicate $O_P : Y^X \rightarrow [0, 1]$, is called fuzzifying pre-openness, is given as: $O_P(f) := \forall U(U \in \tau_P \rightarrow f(U) \in \sigma_P)$. Intuitively, the degree to which f is pre-open is*

$$[O_P(f)] = \bigwedge_{B \subseteq X} \min(1, 1 - \tau_P(U) + \sigma_P(f(U))).$$

Example 3.2. Let (X, τ) and τ_P be defined just as in Example 2.1. We set $Y = X, \sigma = \tau$ and $f = id_X$, then $[O_P(f)] = \bigwedge_{U \subseteq X} \min(1, 1 - \tau_P(U) + \sigma_P(f(U))) = 1$.

Lemma 3.1. Let (X, τ) and (Y, σ) be two fuzzifying topological space. For any $f \in Y^X$,

$$O_P(f) := \forall B (B \in \beta_P^X \rightarrow f(B) \in \sigma_P),$$

where β_P^X is a pre-base of τ_P .

Proof. Clearly, $[O_P(f)] \leq [\forall U (U \in \beta_P^X \rightarrow f(U) \in \sigma_P)]$. Conversely, for any $U \subseteq X$, we are going to prove

$$\min(1, 1 - \tau_P(U) + \sigma_P(f(U))) \geq [\forall V (V \in \beta_P^X \rightarrow f(V) \in \sigma_P)].$$

If $\tau_P(U) \leq \sigma_P(f(U))$, it is hold clearly. Now, assume $\tau_P(U) > \sigma_P(f(U))$. If $\mathfrak{R} \subseteq P(X)$ with $\bigcup \mathfrak{R} = U$, then $\bigcup_{V \in \mathfrak{R}} f(V) = f(\bigcup \mathfrak{R}) = f(U)$. Therefore,

$$\begin{aligned} \tau_P(U) - \sigma_P(f(U)) &= \bigvee_{\mathfrak{R} \subseteq P(X), \bigcup \mathfrak{R} = U} \bigwedge_{V \in \mathfrak{R}} \beta_P^X(V) - \bigvee_{\wp \subseteq P(Y), \bigcup \wp = f(U)} \bigwedge_{W \in \wp} \sigma_P(W) \\ &\leq \bigvee_{\mathfrak{R} \subseteq P(X), \bigcup \mathfrak{R} = U} \bigwedge_{V \in \mathfrak{R}} \beta_P^X(V) - \bigvee_{\mathfrak{R} \subseteq P(X), \bigcup \mathfrak{R} = U} \bigwedge_{V \in \mathfrak{R}} \sigma_P(f(V)) \\ &\leq \bigvee_{\mathfrak{R} \subseteq P(X), \bigcup \mathfrak{R} = U} \bigwedge_{V \in \mathfrak{R}} (\beta_P^X(V) - \sigma_P(f(V))), \\ \min(1, 1 - \tau_P(U) + \sigma_P(f(U))) &\geq \bigvee_{\mathfrak{R} \subseteq P(X), \bigcup \mathfrak{R} = U} \bigwedge_{V \in \mathfrak{R}} \min(1, 1 - \beta_P^X(V) + \sigma_P(f(V))) \\ &\geq [\forall V (V \in \beta_P^X \rightarrow f(V) \in \sigma_P)]. \quad \square \end{aligned}$$

Lemma 3.2. For any family $\{(X_s, \tau_s) : s \in S\}$ of fuzzifying topological spaces.

- (1) $\models (\forall s)(s \in S \rightarrow p_s \in O_P)$;
- (2) $\models (\forall s)(s \in S \rightarrow p_s \in C_P)$.

Proof. (1) For any $t \in S$, we have

$$O_P(p_t) = \bigwedge_{U \in P(\prod_{s \in S} X_s)} \min(1, 1 - \left(\prod_{s \in S} (\tau_P)_s \right) (U) + (\tau_P)_t (p_t(U))).$$

Then, it suffices to show that for any $U \in P(\prod_{s \in S} X_s)$, we have

$$(\tau_P)_t(p_t(U)) \geq \left(\prod_{s \in S} (\tau_P)_s \right) (U).$$

Assume

$$\left(\prod_{s \in S} (\tau_P)_s \right) (U) = \bigvee_{\substack{\cup B_\lambda = U \\ \lambda \in \Lambda}} \bigwedge_{\lambda \in \Lambda} \bigvee_{\Phi_\lambda \sqsubset \varphi, \cap \Phi_\lambda = B_\lambda} \bigwedge_{s \in S(\Phi_\lambda)} (\tau_P)_s(V_s) > \mu$$

where

$$\Phi_\lambda = \{p_s^{-1}(V_s) : s \in S(\Phi_\lambda)\} (\lambda \in \Lambda).$$

Hence, there exists $\{B_\lambda : \lambda \in \Lambda\} \subseteq P(\prod_{s \in S} X_s)$ such that $\bigcup_{\lambda \in \Lambda} B_\lambda = U$ and furthermore, for any $\lambda \in \Lambda$, there exists $\Phi_\lambda \sqsubset \varphi$ such that $\cap \Phi_\lambda = B_\lambda$ and $\bigcap_{s \in S(\Phi_\lambda)} p_s^{-1}(V_s) = B_\lambda$, where for any $s \in S(\Phi_\lambda)$ we have $(\tau_P)_s(V_s) > \mu$.

Thus,

$$p_t(U) = p_t\left(\bigcup_{\lambda \in \Lambda} \bigcap_{s \in S(\Phi_\lambda)} p_s^{-1}(V_s)\right).$$

(1) If for any $\lambda \in \Lambda$, $\bigcap_{s \in S(\Phi_\lambda)} p_s^{-1}(V_s) = \phi$, then $U = \phi, p_t(U) = \phi$ and

$$(\tau_P)_t(p_t(U)) = 1. \text{ Therefore, } (\tau_P)_t(p_t(U)) \geq \left(\prod_{s \in S} (\tau_P)_s \right) (U).$$

(2) If there exists $\lambda_0 \in \Lambda$, such that $\phi \neq \bigcap_{s \in S(\Phi_{\lambda_0})} p_s^{-1}(V_s) = B_{\lambda_0}$,

(i) If $t \notin S(\Phi_{\lambda_0})$, i.e., $t \in S - S(\Phi_{\lambda_0})$, $p_t(B_{\lambda_0}) = X_t$. Therefore, $(\tau_P)_t(p_t(B_{\lambda_0})) = (\tau_P)_t(X_t) = 1$.

(ii) If $t \in S(\Phi_{\lambda_0})$, then $p_t(B_{\lambda_0}) = V_t \subseteq X_t$. Thus,

$$\begin{aligned} p_t(U) &= p_t\left(\left(\bigcup_{t \in S(\Phi_{\lambda_0})} B_{\lambda_0}\right) \cup \left(\bigcup_{t \notin S(\Phi_{\lambda_0})} B_{\lambda_0}\right)\right) \\ &= \left(\bigcup_{t \in S(\Phi_{\lambda_0})} p_t(B_{\lambda_0})\right) \cup \left(\bigcup_{t \notin S(\Phi_{\lambda_0})} p_t(B_{\lambda_0})\right) = V_t \cup X_t = X_t. \end{aligned}$$

Hence,

$$(\tau_P)_t(p_t(U)) = (\tau_P)_t(X_t) = 1 \text{ or } (\tau_P)_t(p_t(U)) = (\tau_P)_t(V_t) > \lambda.$$

Therefore, $(\tau_P)_t(p_t(U)) \geq \left(\prod_{s \in S} (\tau_P)_s \right)(U)$. Thus $O_P(p_t) = 1$.

(2) From Lemma 3.1 in [32], we have $(\forall s)(s \in S \rightarrow p_s \in C)$. Furthermore, for any two fuzzifying topological spaces (X, τ) and (Y, σ) and $f \in Y^X$, we have $C(f) \leq C_P(f)$ (Theorem 6.3 in [2]). Therefore, $\models (\forall s)(s \in S \rightarrow p_s \in C_P)$. \square

Theorem 3.2. *Let $\{(X_s, \tau_s) : s \in S\}$ be the family of fuzzifying topological spaces, then*

$$\begin{aligned} & \models \exists U(U \subseteq \prod_{s \in S} X_s \wedge \Gamma_P(U, \tau/U) \wedge \exists x(x \in \text{Int}_P(U))) \\ & \rightarrow \exists T(T \sqsubset S \wedge \forall t(t \in S - T \wedge \Gamma_P(X_t, \tau_t))). \end{aligned}$$

Proof. It suffices to show that

$$\begin{aligned} & \bigvee_{U \in P(\prod_{s \in S} X_s)} \left(\Gamma_P(U, \tau/U) \wedge \bigvee_{x \in \prod_{s \in S} X_s} N_x^P(U) \right) \\ & \leq \bigvee_{T \sqsubset S} \bigwedge_{t \in S - T} \Gamma_P(X_t, \tau_t). \end{aligned}$$

Indeed, if

$$\bigvee_{U \in P(\prod_{s \in S} X_s)} \left(\Gamma_P(U, \tau/U) \wedge \bigvee_{x \in \prod_{s \in S} X_s} N_x^P(U) \right) > \mu > 0,$$

then there exists $U \in P(\prod_{s \in S} X_s)$ such that $\Gamma_P(U, \tau/U) > \mu$ and

$$\bigvee_{x \in \prod_{s \in S} X_s} N_x^P(U) > \mu, \text{ where}$$

$$N_x^P(U) = \bigvee_{x \in V \subseteq U} \left(\prod_{s \in S} (\tau_P)_s \right)(V).$$

Furthermore, there exists V such that $x \in V \subseteq U$ and $\left(\prod_{s \in S} (\tau_P)_s\right)(V) > \mu$. Since β_P is a pre-base of $\prod_{s \in S} (\tau_P)_s$,

$$\begin{aligned} \left(\prod_{s \in S} (\tau_P)_s\right)(V) &= \bigcup_{\lambda \in \Lambda} \bigvee_{B_\lambda = V} \bigwedge_{\lambda \in \Lambda} \beta_P(B_\lambda) \\ &= \bigcup_{\lambda \in \Lambda} \bigvee_{B_\lambda = V} \bigwedge_{\lambda \in \Lambda} \bigvee_{\Phi_\lambda \sqsubset \varphi, \cap \Phi_\lambda = B_\lambda} \bigwedge_{s \in S(\Phi_\lambda)} (\tau_P)_s(V_s) > \mu, \end{aligned}$$

where $\Phi_\lambda = \{p_s^{-1}(V_s) : s \in S(\Phi_\lambda)\} (\lambda \in \Lambda)$.

Hence there exists $\{B_\lambda : \lambda \in \Lambda\} \subseteq P(\prod_{s \in S} X_s)$ such that $\cup_{\lambda \in \Lambda} B_\lambda = V$. Furthermore, for any $\lambda \in \Lambda$, there exists $\Phi_\lambda \sqsubset \varphi$ such that $\cap \Phi_\lambda = B_\lambda$ and for any $s \in S(\Phi_\lambda)$, we have $(\tau_P)_s(V_s) > \mu$. Since $x \in V$, there exists B_{λ_x} such that $x \in B_{\lambda_x} \subseteq V \subseteq U$. Hence there exists $\Phi_{\lambda_x} \sqsubset \varphi$ such that $\cap \Phi_{\lambda_x} = B_{\lambda_x}$ and $\bigcap_{s \in S(\Phi_{\lambda_x})} p_s^{-1}(V_s) = B_{\lambda_x} \subseteq \prod_{s \in S} X_s$ and for any $s \in S(\Phi_{\lambda_x})$, we have $(\tau_P)_s(V_s) > 1 - \mu$. By $\bigcap_{s \in S(\Phi_{\lambda_x})} p_s^{-1}(V_s) = B_{\lambda_x}$, we have $P_\delta(B_{\lambda_x}) = V_\delta \subseteq X_\delta$, if $\delta \in S(\Phi_{\lambda_x})$; $P_\delta(B_{\lambda_x}) = X_\delta$, if $\delta \in S - S(\Phi_{\lambda_x})$. Since $B_{\lambda_x} \subseteq U$, for any $\delta \in S - S(\Phi_{\lambda_x})$, we have $P_\delta(U) \supseteq P_\delta(B_{\lambda_x}) = X_\delta$ and $P_\delta(U) = X_\delta$. On the other hand, since for any $s \in S$ and $U_s \in P(X_s)$, $\left(\prod_{t \in S} (\tau_P)_t\right)(p_s^{-1}(U_s)) \geq (\tau_P)_s(U_s)$, we have, for any $s \in S$,

$$I_P(p_s) = \bigwedge_{U_s \in P(X_s)} \min \left(1, 1 - (\tau_P)_s(U_s) + \prod_{t \in S} (\tau_P)_t(p_s^{-1}(U_s)) \right) = 1.$$

Furthermore, since by Theorem 9 in [1], we have

$$\models \Gamma_P(X, \tau) \otimes I_P(f) \rightarrow \Gamma_P(f(X)),$$

then

$$\Gamma_P(U, \tau/U) = \Gamma_P(U, \tau/U) \otimes I_P(p_s) \leq \Gamma_P(P_\delta(U), \tau_\delta) = \Gamma_P(X_\delta, \tau_\delta).$$

Therefore,

$$\bigvee_{T \sqsubset S} \bigwedge_{t \in S-T} \Gamma_P(X_t, \tau_t) \geq \bigwedge_{\delta \in S-S(\Phi_\lambda)} \Gamma_P(X_\delta, \tau_\delta) \geq \Gamma_P(U, \tau/U) > \mu.$$

□

The above theorem is a generalization of the following corollary.

Corollary 3.2. *If there exists a coordinate pre-neighborhood strong compact subset U of some point $x \in X$ of the product space, then all except a finite number of coordinate spaces are strong compact.*

Lemma 3.3. *For any fuzzifying topological space (X, τ) , $A \subseteq X$,
 $\models T_2^P(X, \tau) \rightarrow T_2^P(A, \tau/A)$.*

Proof.

$$\begin{aligned} [T_2^P(X, \tau)] &= \bigwedge_{x,y \in X, x \neq y} \bigvee_{U,V \in P(X), U \cap V = \phi} (N_x^P(U), N_y^P(V)) \\ &\leq \bigwedge_{x,y \in X, x \neq y} \bigvee_{(U \cap A) \cap (V \cap A) = \phi} (N_x^{PA}(U \cap A), N_y^{PA}(V \cap A)) \\ &\leq \bigwedge_{x,y \in A, x \neq y} \bigvee_{U' \cap V' = \phi, U', V' \in P(A)} (N_x^{PA}(U'), N_y^{PA}(V')) \\ &= T_2^P(A, \tau/A), \end{aligned}$$

where

$$N_x^{PA}(U) = \bigvee_{x \in C \subseteq U} \tau_{P/A}(C) \text{ and } \tau_{P/A}(B) = \bigvee_{B=V \cap A} \tau_P(V).$$

□

Lemma 3.4. *For any fuzzifying P -topological space (X, τ) ,
 $\models T_2^P(X, \tau) \otimes \Gamma_P(X, \tau) \rightarrow T_4^P(X, \tau)$.
 For the definition of $T_4^P(X, \tau)$ see [3, Definition 3.1].*

Proof. If $[T_2^P(X, \tau) \otimes \Gamma_P(X, \tau)] = 0$, then the result holds. Now, suppose that $[T_2^P(X, \tau) \otimes \Gamma_P(X, \tau)] > \lambda > 0$. Then $T_2^P(X, \tau) + \Gamma_P(X, \tau) - 1 > \lambda > 0$. Therefore, from Theorem 10 in [1]
 $T_2^P(X, \tau) \otimes (\Gamma_P(A) \wedge \Gamma_P(B)) \wedge (A \cap B = \phi) \models^{ws} T_2^P(X, \tau) \rightarrow (\exists U)(\exists V)((U \in \tau_P) \wedge (V \in \tau_P) \wedge (A \subseteq U) \wedge (B \subseteq V) \wedge (A \cap B = \phi))$. Then for any $A, B \subseteq X, A \cap B = \phi$,

$$T_2^P(X, \tau) \otimes (\Gamma_P(A) \wedge \Gamma_P(B)) \leq \bigvee_{U \cap V = \phi, A \subseteq U, B \subseteq V} \min(\tau_P(U), \tau_P(V))$$

or equivalently

$$T_2^P(X, \tau) \leq \Gamma_P(A) \wedge \Gamma_P(B) \rightarrow \bigvee_{U \cap V = \phi, A \subseteq U, B \subseteq V} \min(\tau_P(U), \tau_P(V))$$

Hence, for any $A, B \subseteq X, A \cap B = \phi$,

$$1 - [\Gamma_P(A) \wedge \Gamma_P(B)] + \bigvee_{U \cap V = \phi, A \subseteq U, B \subseteq V} \min(\tau_P(U), \tau_P(V)) + \Gamma_P(X, \tau) - 1 > \lambda.$$

From Theorem 7 in [1] we have

$$\models \Gamma_P(X, \tau) \otimes A \in \mathcal{F}_P \rightarrow \Gamma_P(A).$$

Then

$$\begin{aligned} \Gamma_P(X, \tau) + [\tau_P(A^c) \wedge \tau_P(B^c)] - 1 &= (\Gamma_P(X, \tau) + \tau_P(A^c) - 1) \wedge (\Gamma_P(X, \tau) + \tau_P(B^c) - 1) \\ &\leq (\Gamma_P(X, \tau) \otimes \tau_P(A^c)) \wedge (\Gamma_P(X, \tau) \otimes \tau_P(B^c)) \\ &\leq [\Gamma_P(A) \wedge \Gamma_P(B)]. \end{aligned}$$

Thus,

$$\Gamma_P(X, \tau) - [\Gamma_P(A) \wedge \Gamma_P(B)] - 1 \leq -[\tau_P(A^c) \wedge \tau_P(B^c)].$$

So,

$$1 - [\tau_P(A^c) \wedge \tau_P(B^c)] + \bigvee_{U \cap V = \phi, A \subseteq U, B \subseteq V} \min(\tau_P(U), \tau_P(V)) > \lambda,$$

i.e.,

$$\begin{aligned} T_4^P(X, \tau) &= \bigwedge_{A \cap B = \phi} \min(1, 1 - [\tau_P(A^c) \wedge \tau_P(B^c)]) \\ &+ \bigvee_{U \cap V = \phi, A \subseteq U, B \subseteq V} \min(\tau_P(U), \tau_P(V)) > \lambda. \end{aligned}$$

□

The above lemma is a generalization of the following corollary.

Corollary 3.3. *Every strong compact pre-Hausdorff topological space is pre-normal.*

Lemma 3.5. *For any fuzzifying P -topological space (X, τ) , $\models T_2^P(X, \tau) \otimes \Gamma_P(X, \tau) \rightarrow T_3^P(X, \tau)$. For the definition of $T_3^P(X, \tau)$ see [3, Definition 3.1].*

Proof. Immediate, set $A = \{x\}$ in the above lemma. \square

The above lemma is a generalization of the following corollary.

Corollary 3.4. *Every strong compact pre-Hausdorff topological space is pre-regular.*

Theorem 3.3. *For any fuzzifying topological space (X, τ) and $A \subseteq X$, $\models T_2^P(X, \tau) \otimes \Gamma_P(A) \rightarrow A \in \mathcal{F}_P$.*

Proof. For any $\{x\} \subset A^c$, we have $\{x\} \cap A = \phi$ and $\Gamma_P(\{x\}) = 1$. By Theorem 10 in [1]

$$[T_2^P(X, \tau) \otimes (\Gamma_P(A) \wedge \Gamma_P(\{x\}))] \leq \bigvee_{G \cap H_x = \phi, A \subseteq G, x \in H_x} \min(\tau_P(G), \tau_P(H_x)).$$

Assume

$$\beta_x = \{H_x : A \cap H_x = \phi, x \in H_x\}, \quad \bigcup_{x \in X \setminus A} f(x) \supseteq A^c$$

and

$$\bigcup_{x \in A^c} f(x) \cap A = \bigcup_{x \in A^c} (f(x) \cap A) = \phi.$$

So, $\bigcup_{x \in A^c} f(x) = A^c$.

Therefore,

$$\begin{aligned} [T_2^P(X, \tau) \otimes \Gamma_P(A)] &\leq \bigvee_{G \cap H_x = \phi, A \subseteq G, x \in H_x} \tau_P(H_x) \\ &\leq \bigwedge_{x \in A^c} \bigvee_{A \cap H_x = \phi, x \in H_x} \tau_P(H_x) \\ &= \bigvee_{f \in \prod_{x \in A^c} \beta_x} \bigwedge_{x \in A^c} \tau_P(f(x)) \\ &\leq \bigvee_{f \in \prod_{x \in A^c} \beta_x} \tau_P\left(\bigcup_{x \in A^c} f(x)\right) \\ &= \bigvee_{f \in \prod_{x \in X \setminus A} \beta_x} \tau_P(A^c) = \mathcal{F}_P(A). \end{aligned}$$

\square

The above theorem is a generalization of the following corollary.

Corollary 3.5. *Strong compact subspace of a pre-Hausdorff topological space is pre-closed.*

4. Fuzzifying locally strong compactness

Definition 4.1. Let Ω be a class of fuzzifying topological spaces. A unary fuzzy predicate $L_P C : \Omega \rightarrow [0, 1]$, called *fuzzifying locally strong compactness*, is given as follows:

$$(X, \tau) \in L_P C := (\forall x)(\exists B)((x \in \text{Int}_P(B) \otimes \Gamma_P(B, \tau/B)).$$

Since $[x \in \text{Int}_P(X)] = N_x^P(X) = 1$, then $L_P C(X, \tau) \geq \Gamma_P(X, \tau)$. Therefore, $\models (X, \tau) \in \Gamma_P \rightarrow (X, \tau) \in L_P C$.

Also, since $\models (X, \tau) \in \Gamma \rightarrow (X, \tau) \in LC$ [21] and $\models (X, \tau) \in \Gamma_P \rightarrow (X, \tau) \in \Gamma$ [1], $\models (X, \tau) \in \Gamma_P \rightarrow (X, \tau) \in LC$.

Theorem 4.1. For any fuzzifying topological space (X, τ) and $A \subseteq X$, $\models (X, \tau) \in L_P C \otimes A \in F_P \rightarrow (A, \tau/A) \in L_P C$.

Proof. We have

$$L_P C(X, \tau) = \bigwedge_{x \in X} \bigvee_{B \subseteq X} \max(0, N_x^{P^X}(B) + \Gamma_P(B, \tau/B) - 1)$$

and

$$L_P C(A, \tau/A) = \bigwedge_{x \in A} \bigvee_{G \subseteq A} \max(0, N_x^{P^A}(G) + \Gamma_P(G, (\tau/A)/G) - 1).$$

Now, suppose that $[(X, \tau) \in L_P C \otimes A \in F_P] > \lambda > 0$. Then for any $x \in A$, there exists $B \subseteq X$ such that

$$(4.1) \quad N_x^{P^X}(B) + \Gamma_P(B, \tau/B) + \tau_P(X - A) - 2 > \lambda$$

Set $E = A \cap B \in P(A)$. Then,

$$N_x^{P^A}(E) = \bigvee_{E=C \cap B} N_x^{P^X}(C) \geq N_x^{P^X}(B)$$

and for any $U \in P(E)$, we have

$$\begin{aligned} (\tau_P/A)_{P/E}(U) &= \bigvee_{U=C \cap E} \tau_P/A(C) \\ &= \bigvee_{U=C \cap E} \bigvee_{C=D \cap A} \tau_P(D) \\ &= \bigvee_{U=D \cap A \cap E} \tau_P(D) = \bigvee_{U=D \cap E} \tau_P(D). \end{aligned}$$

Similarly,

$$(\tau_P/B)_P/E(U) = \bigvee_{U=D \cap E} \tau_P(D).$$

Thus, $(\tau_P/B)_P/E = (\tau_P/A)_P/E$ and $\Gamma_P(E, (\tau/A)/E) = \Gamma_P(E, (\tau/B)/E)$. Furthermore,

$$\begin{aligned} [E \in F_P/B] &= \tau_P/B(B - E) = \tau_P/B(B \cap E^c) \\ &= \bigvee_{B \cap E^c = B \cap D} \tau_P(D) \\ &\geq \tau_P(X - A) = F_P(A). \end{aligned}$$

Since $\models (X, \tau) \in \Gamma_P \otimes A \in F_P \rightarrow (A, \tau/A) \in \Gamma_P$ (see [1], Theorem 7], from (4.1) we have for any $x \in A$ that

$$\begin{aligned} &\bigvee_{G \subseteq A} \max(0, N_x^{PA}(G) + \Gamma_P(G, (\tau/A)/G) - 1) \\ &\geq N_x^{PA}(E) + \Gamma_P(E, (\tau/A)/E) - 1 \\ &= N_x^{PA}(E) + \Gamma_P(E, (\tau/B)/E) - 1 \\ &\geq N_x^{PX}(B) + [\Gamma_P(B, \tau/B) \otimes E \in F_P/B] - 1 \\ &\geq N_x^{PX}(B) + \Gamma_P(B, \tau/B) + [E \in F_P/B] - 2 \\ &\geq N_x^{PX}(B) + \Gamma_P(B, \tau/B) + [A \in F_P] - 2 > \lambda. \end{aligned}$$

Therefore,

$$L_P C(A, \tau/A) = \bigwedge_{x \in A} \bigvee_{G \subseteq A} \max(0, N_x^{PA}(G) + \Gamma_P(G, (\tau/A)/G) - 1) > \lambda.$$

Hence, $[(X, \tau) \in L_P C \otimes A \in F_P] \leq L_P C(A, \tau/A)$. \square

As a crisp result of the above theorem we have the following corollary.

Corollary 4.1. *Let A be a pre-closed subset of locally strong compact space (X, τ) . Then A with the relative topology τ/A is locally strong compact.*

The following theorem is a generalization of the statement "If X is a pre-Hausdorff topological space and A is a pre-dense locally strong compact subspace, then A is pre-open", where A is a pre-dense in a topological space X if and only if the pre-closure of A is X .

Theorem 4.2. *For any fuzzifying P -topological space (X, τ) and $A \subseteq X$,*

$$\models T_2^P(X, \tau) \otimes L_P C(A, \tau/A) \otimes (Cl_P(A) \equiv X) \rightarrow A \in \tau_P.$$

Proof. Assume

$$[T_2^P(X, \tau) \otimes L_P C(A, \tau/A) \otimes (Cl_P(A) \equiv X)] > \lambda > 0.$$

Then,

$$L_P C(A, \tau/A) > \lambda - [T_2^P(X, \tau) \otimes (Cl_P(A) \equiv X)] + 1 = \lambda' > \lambda,$$

i. e.,

$$\bigwedge_{x \in A} \bigvee_{B \subseteq A} \max(0, N_x^{PA}(B) + \Gamma_P(B, (\tau/A)/B) - 1) > \lambda'.$$

Thus, for any $x \in A$, there exists $B_x \subseteq A$ such that

$$N_x^{PA}(B_x) + \Gamma_P(B_x, (\tau/A)/B_x) - 1 > \lambda'.$$

i.e.,

$$\bigvee_{H \cap A = B_x} \bigvee_{x \in K \subseteq H} \tau_P(K) + \Gamma_P(B_x, (\tau/A)/B_x) - 1 > \lambda'.$$

Hence, there exists K_x such that

$$K_x \cap A = B_x, \tau_P(K_x) + \Gamma_P(B_x, (\tau/A)/B_x) - 1 > \lambda'.$$

Therefore, $\tau_P(K_x) > \lambda'$.

(1) If for any $x \in A$ there exists K_x such that

$$x \in K_x \subseteq B_x \subseteq A, \text{ then } \bigcup_{x \in A} K_x = A$$

and

$$\tau_P(A) = \tau_P\left(\bigcup_{x \in A} K_x\right) \geq \bigwedge_{x \in A} \tau_P(K_x) \geq \lambda' > \lambda.$$

(2) If there exists $x_o \in A$ such that

$$K_{x_o} \cap (B_{x_o}^c) \neq \phi, \tau_P(K_{x_o}) + \Gamma_P(B_{x_o}, (\tau/A)/B_{x_o}) - 1 > \lambda'.$$

From the hypothesis

$$[T_2^P(X, \tau) \otimes L_P C(A, \tau/A) \otimes (Cl_P(A) \equiv X)] > \lambda > 0,$$

we have $[T_2^P(X, \tau) \otimes (Cl_P(A) \equiv X)] \neq 0$. So

$$\tau_P(K_{x_o}) + \Gamma_P(B_{x_o}, (\tau/A)/B_{x_o}) - 1 + [T_2^P(X, \tau) \otimes (Cl_P(A) \equiv X)] - 1 > 0.$$

Therefore,

$$\tau_P(K_{x_0}) + \Gamma_P(B_{x_0}, (\tau/A)/B_{x_0}) - 1 + T_2^P(X, \tau) + [(Cl_P(A) \equiv X)] - 1 - 1 > \lambda.$$

Since

$$\begin{aligned} (\tau_P/A)_{P/B_{x_0}}(U) &= \bigvee_{U=C \cap B_{x_0}} \tau_P/A(C) \\ &= \bigvee_{U=C \cap B_{x_0}} \bigvee_{C=D \cap A} \tau_P(D) \\ &= \bigvee_{U=D \cap B_{x_0}} \tau_P(D) = \tau_{P/B_{x_0}}(U), \\ \Gamma_P(B_{x_0}, (\tau/A)/B_{x_0}) &= \Gamma_P(B_{x_0}, \tau/B_{x_0}). \end{aligned}$$

From Theorem 3.3 we have

$$\begin{aligned} \tau_P(B_{x_0}^c) &\geq T_2^P(X, \tau) \otimes \Gamma_P(B_{x_0}, \tau/B_{x_0}) \\ &\geq T_2^P(X, \tau) + \Gamma_P(B_{x_0}, \tau/B_{x_0}) - 1. \end{aligned}$$

Hence,

$$\tau_P(K_{x_0}) + \tau_P(B_{x_0}^c) + [Cl_P(A) \equiv X] - 2 > \lambda.$$

Now, for any $y \in A^c$ we have

$$[Cl_P(A) \equiv X] = \bigwedge_{x \in X} (1 - N_x^{P^X}(A^c)) \leq 1 - N_y^{P^X}(A^c).$$

Since (X, τ) is a fuzzifying P -topological space,

$$\begin{aligned} \tau_P(K_{x_0}) + \tau_P(B_{x_0}^c) - 1 &\leq \tau_P(K_{x_0}) \otimes \tau_P(B_{x_0}^c) \\ &\leq \tau_P(K_{x_0}) \wedge \tau_P(B_{x_0}^c) \\ &\leq \tau_P(K_{x_0} \cap B_{x_0}^c) \\ &\leq N_y^{P^X}(K_{x_0} \cap B_{x_0}^c) \leq N_y^{P^X}(A^c), \end{aligned}$$

where

$$\begin{aligned} y \in K_{x_0} \cap B_{x_0}^c &\subseteq H_{x_0} \cap (H_{x_0} \cap A)^c = H_{x_0} \cap (H_{x_0}^c \cup A^c) \\ &= H_{x_0} \cap A^c \subseteq A^c. \end{aligned}$$

Therefore,

$$\begin{aligned} 0 &< \lambda < \tau_P(K_{x_0}) + \tau_P(B_{x_0}^c) + [Cl_P(A) \equiv X] - 2 \\ &= \tau_P(K_{x_0}) + \tau_P(B_{x_0}^c) - 1 + [Cl_P(A) \equiv X] - 1 \\ &\leq N_y^{P^X}(A^c) + 1 - N_y^{P^X}(A^c) - 1 = 0, \end{aligned}$$

a contradiction. So, case (2) does not hold. We complete the proof. \square

Theorem 4.3. For any fuzzifying P -topological space (X, τ) ,

$$\begin{aligned} & \models T_2^P(X, \tau) \otimes (L_P C(X, \tau))^2 \\ & \rightarrow \forall x \forall U (U \in N_x^{P^X} \rightarrow \exists V (V \in N_x^{P^X} \wedge Cl_P(V) \subseteq U \wedge \Gamma_P(V))), \end{aligned}$$

where $(L_P C(X, \tau))^2 := L_P C(X, \tau) \otimes L_P C(X, \tau)$.

Proof. We need to show that for any x and U , $x \in U$,

$$\begin{aligned} & T_2^P(X, \tau) \otimes (L_P C(X, \tau))^2 \otimes N_x^{P^X}(U) \\ & \leq \bigvee_{V \subseteq X} (N_x^{P^X}(V) \wedge \bigwedge_{y \in U^c} N_x^{P^X}(V^c) \wedge \Gamma_P(V, \tau/V)). \end{aligned}$$

Assume that $T_2^P(X, \tau) \otimes (L_P C(X, \tau))^2 \otimes N_x^{P^X}(U) > \lambda > 0$. Then, for any $x \in X$ there exists C such that

$$(4.2) \quad T_2^P(X, \tau) + N_x^{P^X}(C) + (L_P C(X, \tau))^2 + N_x^{P^X}(U) - 3 > \lambda.$$

Since (X, τ) is fuzzifying P -topological space,

$$\begin{aligned} & N_x^{P^X}(C) + N_x^{P^X}(U) - 1 \leq N_x^{P^X}(C) \otimes N_x^{P^X}(U) \\ & \leq N_x^{P^X}(C) \wedge N_x^{P^X}(U) \\ & \leq N_x^{P^X}(C \cap U) = \bigvee_{x \in W \subseteq C \cap U} \tau_P(W). \end{aligned}$$

Therefore, there exists W such that $x \in W \subseteq C \cap U$, and $T_2^P(X, \tau) + (L_P C(X, \tau))^2 + \tau_P(W) - 2 > \lambda$. By Lemmas 3.3 and 3.5 we have $T_2^P(X, \tau) \leq T_2^P(C, \tau/C)$ and

$$T_2^P(C, \tau/C) + \Gamma_P(C, \tau/C) - 1 \leq T_2^P(C, \tau/C) \otimes \Gamma_P(C, \tau/C) \leq T_3^P(C, \tau/C).$$

Thus, $T_3^P(X, \tau) + \Gamma_P(C, \tau/C) + \tau_P(W) - 2 > \lambda$. Since for any $x \in W \subseteq U$, we have

$$T_3^P(C, \tau/C) \leq 1 - \tau_{P/C}(W) + \bigvee_{G \subseteq C} \left((N_x^{P^C}(G) \wedge \bigwedge_{y \in C-W} N_y^{P^C}(C-G)) \right),$$

so there exists $G, x \in G \subseteq W$ such that

$$\begin{aligned} & \left(N_x^{PC}(G) \wedge \bigwedge_{y \in C-W} N_y^{PC}(C-G) \right) \\ & \geq T_3^P(C, \tau/C) + \tau_P/C(W) - 1 \geq T_3^P(C, \tau/C) + \tau_P(W) - 1 \end{aligned}$$

and

$$\left(N_x^{PC}(G) \wedge \bigwedge_{y \in C-W} N_y^{PC}(C-G) \right) + \Gamma_P(C, \tau/C) - 1 > \lambda.$$

Thus,

$$N_x^{PC}(G) = \bigvee_{D \cap C = G} N_x^{PX}(D) = N_x^{PX}(G \cup C^c) > \lambda' = \lambda + 1 - \Gamma_P(C, \tau/C) \geq \lambda.$$

Furthermore, for any $y \in C - W$,

$$N_y^{PC}(C-G) = \bigvee_{D \cap C = C \cap G^c} N_y^{PX}(G^c \cup C^c) = N_y^{PX}(G^c) > \lambda'$$

and

$$N_x^{PX}(G) = N_x^{PX}((G \cup C^c) \cap C) \geq N_x^{PX}(G^c) \wedge N_x^{PX}(C) > \lambda'.$$

Since $N_y^{PX}(G^c) = \bigvee_{x \in B^c \subseteq G^c} \tau_P(B^c) > \lambda'$, for any $y \in C - W$, there exists B_y^c such that $y \in B_y^c \subseteq G^c$ and $\tau_P(B_y^c) > \lambda'$. Set $B^c = \bigcup_{y \in C-W} B_y^c$. Then $C - W \subseteq B^c \subseteq G^c$ and

$$\tau_P(B^c) \geq \bigwedge_{y \in C-W} \tau_P(B_y^c) \geq \lambda'.$$

Again, set $V = B \cap C$, then $V \subseteq (C - W)^c \cap C = (C^c \cup W) \cap C = C \cap W = W \subseteq U \cap C$ and $V^c = B^c \cup C^c$. Since (X, τ) is fuzzifying P -topological space,

$$\begin{aligned} (4.3) \quad N_x^{PX}(V) = N_x^{PX}(B \cap C) & \geq N_x^{PX}(B) \wedge N_x^{PX}(C) \\ & \geq N_x^{PX}(G) \wedge N_x^{PX}(C) > \lambda. \end{aligned}$$

By (4.2) and Theorem 3.3,

$$\tau_P(C^c) \geq T_2^P(X, \tau) \otimes \Gamma_P(C, \tau/C) \geq T_2^P(X, \tau) + \Gamma_P(C, \tau/C) - 1 \geq \lambda'.$$

So

$$\tau_P(V^c) = \tau_P(B^c \cup C^c) \geq \tau_P(B^c) \wedge \tau_P(C^c) \geq \lambda',$$

i.e., $\tau_P(V^c) + \Gamma_P(C, \tau/C) - 1 \geq \lambda$ and

$$\begin{aligned} \Gamma_P(V, \tau/V) &= \Gamma_P(V, (\tau/C)/V) \\ &\geq \tau_P/C(C - V) + \Gamma_P(C, \tau/C) - 1 \\ (4.4) \quad &\geq \tau_P(V^c) + \Gamma_P(C, \tau/C) - 1 \geq \lambda \end{aligned}$$

Finally,

$$(4.5) \quad \bigwedge_{y \in U^c} N_y^{PX}(V^c) \geq \bigwedge_{y \in V^c} N_y^{PX}(V^c) = \tau_P(V^c) \geq \lambda$$

Thus, by (4.3), (4.4) and (4.5), for any $x \in U$, there exists $V \subseteq U$ such that $N_x^{PX}(V) > \lambda$, $\bigwedge_{y \in U^c} N_y^{PX}(V^c) \geq \lambda$ and $\Gamma_P(V, \tau/V) \geq \lambda$. So $\bigvee_{V \subseteq X} (N_x^{PX}(V) \wedge \bigwedge_{y \in U^c} N_y^{PX}(V^c) \wedge \Gamma_P(V, \tau/V)) \geq \lambda$. \square

Theorem 4.4. For any fuzzifying P -topological space (X, τ) ,

$$\models T_2^P(X, \tau) \otimes (L_PC(X, \tau))^2 \rightarrow T_3^P(X, \tau).$$

Proof. By Theorem 4.3, for any $x \in U$, we have

$$\begin{aligned} &\bigvee_{x \in V \subseteq U} (N_x^{PX}(V) \wedge \bigwedge_{y \in U^c} N_y^{PX}(V^c)) \\ &\geq [T_2^P(X, \tau) \otimes (\Gamma_P(C, \tau/C))^2 \otimes N_x^{PX}(U)]. \end{aligned}$$

Thus,

$$\begin{aligned} 1 - N_x^{PX}(U) + \bigvee_{x \in V \subseteq U} (N_x^{PX}(V) \wedge \bigwedge_{y \in U^c} N_y^{PX}(V^c)) \\ \geq [T_2^P(X, \tau) \otimes (\Gamma_P(C, \tau/C))^2], \end{aligned}$$

i.e.,

$$[T_3^P(X, \tau)] \geq [T_2^P(X, \tau) \otimes (\Gamma_P(C, \tau/C))^2].$$

\square

Theorem 4.5. For any fuzzifying P -topological space (X, τ) ,

$$\begin{aligned} \models T_3^P(X, \tau) \otimes L_PC(X, \tau) &\rightarrow \forall A \forall U (U \in N_A^{PX} \otimes \Gamma_P(A, \tau/A) \\ &\rightarrow \exists V (V \subseteq U \wedge U \in N_A^{PX} \wedge \tau_P(V^c) \wedge \Gamma_P(V, \tau/V))), \end{aligned}$$

where

$$U \in N_A^{PX} := (\forall x)(x \in A \wedge U \in N_x^{PX}).$$

Proof. We only need to show that for any $A, U \in P(X)$,

$$\begin{aligned} & [T_3^P(X, \tau) \otimes L_P C(X, \tau) \otimes \Gamma_P(A, \tau/A) \otimes N_A^{PX}(U)] \\ & \leq \bigvee_{V \subseteq U} (N_A^{PX}(V) \wedge \tau_P(V^c) \wedge \Gamma_P(V, \tau/V)). \end{aligned}$$

Indeed, if

$$[T_3^P(X, \tau) \otimes L_P C(X, \tau) \otimes \Gamma_P(A, \tau/A) \otimes N_A^{PX}(U)] > \lambda > 0,$$

then for any $x \in A$, there exists $C \in P(X)$ such that

$$[T_3^P(X, \tau) \otimes N_x^{PX}(C) \otimes \Gamma_P(C, \tau/C) \otimes \Gamma_P(A, \tau/A) \otimes N_A^{PX}(U)] > \lambda.$$

Since (X, τ) is fuzzifying P -topological space,

$$\begin{aligned} \bigvee_{x \in W \subseteq C \cap U} \tau_P(W) = N_x^{PX}(C \cap U) & \geq N_x^{PX}(C) \wedge N_x^{PX}(U) \\ & \geq N_x^{PX}(C) \wedge N_A^{PX}(U) \\ & \geq N_x^{PX}(C) \otimes N_A^{PX}(U). \end{aligned}$$

Then, there exists W such that $x \in W \subseteq C \cap U$, and

$$[T_3^P(X, \tau) \otimes \tau_P(W) \otimes \Gamma_P(C, \tau/C) \otimes \Gamma_P(A, \tau/A)] > \lambda.$$

Therefore,

$$(4.6)[T_3^P(X, \tau)] + \tau_P(W) - 1 > \lambda + 2 - \Gamma_P(C, \tau/C) - \Gamma_P(A, \tau/A) = \lambda' \geq \lambda.$$

Since for any $x \in W$,

$$[T_3^P(X, \tau)] \leq 1 - \tau_P(W) + \bigvee_{B \subseteq W} (N_x^{PX}(B) \wedge \bigwedge_{y \in W^c} N_y^{PX}(B^c)),$$

we have

$$\bigvee_{B \subseteq W} (N_x^{PX}(B) \wedge \bigwedge_{y \in W^c} N_y^{PX}(B^c)) > \lambda'.$$

Thus, there exists B_x such that $x \in B_x \subseteq W \subseteq C \cap U$ and for any $y \in W^c$, we have

$$N_y^{PX}(B_x^c) > \lambda', \quad N_x^{PX}(B_x) > \lambda'.$$

Since

$$N_y^{PX}(B_x^c) = \bigvee_{x \in G^c \subseteq B_x^c} \tau_P(G^c) > \lambda',$$

then for any $y \in W^c$, there exists G_{xy} such that

$$x \in G_{xy}^c \subseteq B_x^c \text{ and } \tau_P(G_{xy}^c) > \lambda'.$$

Set

$$G_x^c = \bigcup_{y \in W^c} G_{xy}^c,$$

then $W^c \subseteq G_{xy}^c \subseteq B_x^c$ and $\tau_P(G_x^c) \geq \bigwedge_{y \in W^c} \tau_P(G_{xy}^c) \geq \lambda'$. Since $G_x \supseteq B_x$, $N_x^{PX}(G_x) \geq N_x^{PX}(B_x) > \lambda'$, i.e., $\bigvee_{x \in H \subseteq G_x} \tau_P(H) > \lambda'$. Thus, there exists H_x such that $x \in H_x \subseteq G_x$ and $\tau_P(H_x) > \lambda'$. Hence, for any $x \in A$, there exists H_x and G_x such that $x \in H_x \subseteq G_x \subseteq U$, $\tau_P(H_x) > \lambda'$ and $W \supseteq \bigcup_{x \in A} G_x \supseteq \bigcup_{x \in A} H_x \supseteq A$. We define $\mathfrak{R} \in \mathfrak{S}(P(A))$ as follows:

$$\mathfrak{R}(D) = \begin{cases} \bigvee_{H_x \cap A = D} \tau_P(H_x), & \text{there exists } H_x \text{ such that } H_x \cap A = D, \\ 0, & \text{otherwise.} \end{cases}$$

Let $\Gamma_P(A, \tau/A) = \mu > \mu - \epsilon$ ($\epsilon > 0$). Then $1 - K_P(\mathfrak{R}, A) + \bigvee_{\varphi \leq \mathfrak{R}} [K(\mathfrak{R}, A) \otimes FF(\varphi)] > \mu - \epsilon$, where

$$\begin{aligned} [K(\mathfrak{R}, A)] &= \bigwedge_{x \in A} \bigvee_{x \in B} \mathfrak{R}(B) = \bigwedge_{x \in A} \bigvee_{x \in D} \mathfrak{R}(D) \\ &= \bigwedge_{x \in A} \bigvee_{x \in D} \bigvee_{H_{x'} \cap A = D} \tau_P(H_{x'}) \geq \lambda' \end{aligned}$$

and $[\mathfrak{R} \subseteq \tau_P \setminus A] = \bigwedge_{B \subseteq X} \min(1, 1 - \mathfrak{R}(B) + \tau_P \setminus A(B)) = \bigwedge_{B \subseteq X} \min(1, 1 - \bigvee_{H_x \cap A = B} \tau_P(H_x) + \bigvee_{H \cap A = B} \tau_P(H)) = 1$. So, $K_P(\mathfrak{R}, A) = [K(\mathfrak{R}, A)] \geq \lambda'$. By (4.6), $[K(\mathfrak{R}, A) \otimes FF(\varphi)] > \mu - \epsilon - 1 + K_P(\mathfrak{R}, A) \geq \mu - \epsilon - 1 + \lambda' \geq \lambda - \epsilon$.

Thus, $\bigwedge_{x \in A} \bigvee_{x \in E} \mathfrak{R}(E) + 1 - \bigwedge\{\delta : F(\varphi_\delta)\} - 1 > \lambda - \epsilon$, and $\bigwedge_{x \in A} \bigvee_{x \in E} \mathfrak{R}(E) > \lambda - \epsilon + \bigwedge\{\delta : F(\varphi_\delta)\}$.

Hence, there exists $\beta > 0$ such that $F(\wp\beta)$ and $\bigwedge_{x \in A} \bigvee_{x \in D} \mathfrak{R}(D) > \lambda - \epsilon + \beta$. Therefore for any $x \in A$, there exists $D_x \subseteq A$ such that $\wp(D_x) > \lambda - \epsilon + \beta$ and $\bigcup_{x \in A} D_x \supseteq A$. Suitably choose ϵ such that $\lambda - \epsilon > 0$, then $\wp(D_x) > \beta > 0$. Since $\mathfrak{R}(D_x) \geq \wp(D_x) > 0$, $D_x = H_{x'} \cap A$, i.e., $H_{x'} \cap A \in \wp\beta$. By $F(\wp\beta)$, there exists finite $H_{x'_1}, H_{x'_2}, \dots, H_{x'_n}$ such that $\bigcup_{i=1}^n H_{x'_i} \supseteq A$ and $\bigcup_{i=1}^n H_{x'_i} \subseteq \bigcup_{i=1}^n G_{x'_i}$. Set $V = \bigcup_{i=1}^n G_{x'_i}$, and $V^c = \bigcap_{i=1}^n G_{x'_i}^c$, $A \subseteq V \subseteq U$, and $\tau_P(V^c) \geq \bigwedge_{1 \leq i \leq n} \tau_P(G_{x'_i}^c) \geq \lambda' > \lambda$. Since for any $x \in A$, $G_x \subseteq W \subseteq C \cap U \subseteq C$, we have $V = \bigcup_{i=1}^n G_{x'_i} \subseteq W \subseteq C$. Because $\tau_P \setminus C(C - V) = \bigvee_{D \cap C = C \cap V^c} \tau_P(D) \geq \tau_P(V^c) \geq \lambda'$. Thus by (4.6), $\tau_P \setminus C(C - V) + \Gamma_P(C, \tau/C) - 1 > \lambda$. By Theorem 5.1 in [24], $\Gamma_P(V, \tau/V) = \Gamma_P(V, \tau/C/V) \geq [\Gamma_P(C, \tau/C) \otimes \tau_P \setminus C(C - V)] > \lambda$.

Finally, we have for any $x \in A$, $N_x^{PX}(V) = N_x^{PX}(\bigcup_{i=1}^n G_{x'_i}) \geq N_x^{PX}(\bigcup_{i=1}^n H_{x'_i}) \geq \tau_P(\bigcup_{i=1}^n H_{x'_i}) \geq \bigwedge_{1 \leq i \leq n} \tau_P(H_{x'_i}) \geq \lambda' > \lambda$. So, $N_A^{PX}(V) = \bigwedge_{x \in A} N_x^{PX}(V) \geq \lambda$. Therefore, $N_A^{PX}(V) \wedge \tau_P(V^c) \wedge \Gamma_P(V, \tau/V) \geq \lambda$.

Thus, $\bigvee_{V \subseteq U} (N_A^{PX}(V) \wedge \tau_P(V^c) \wedge \Gamma_P(V, \tau/V)) \geq \lambda$. \square

Theorem 4.6. Let (X, τ) and (Y, σ) be two fuzzifying topological space and $f \in Y^X$ be surjective. Then $\models L_PC(X, \tau) \otimes C_P(f) \otimes O(f) \rightarrow LC(Y, \sigma)$. For the definition of $O(f)$, see [32].

Proof. If $[L_PC(X, \tau) \otimes C_P(f) \otimes O(f)] > \lambda > 0$, then for any $x \in X$, there exists $U \subseteq X$, such that $[N_x^{PX}(U) \otimes \Gamma_P(U, \tau/U) \otimes C_P(f) \otimes O(f)] > \lambda$. Since $N_x^{PX}(U) = \bigvee_{x \in V \subseteq U} \tau_P(V)$, so there exists $V' \subseteq X$ such that $x \in V' \subseteq U$ and $[\tau_P(V') \otimes \Gamma_P(U, \tau/U) \otimes C_P(f) \otimes O(f)] > \lambda$. By Theorem 8 in [1], $[\Gamma_P(U, \tau/U) \otimes C_P(f)] \leq [\Gamma(f(U), \sigma/f(U))]$ and $[\tau(V') \otimes O(f)] = \max(0, \tau(V') + O(f) - 1) = \max(0, \tau(V') + \bigwedge_{V \subseteq X} \min(1, 1 - \tau(V) + \sigma(f(V))) - 1) \leq \max(0, \tau(V') + 1 - \tau(V') + \sigma(f(V)) - 1) = \sigma(f(V)) \leq N_{f(x)}^Y(f(V')) \leq N_{f(x)}^Y(f(U))$. Since f is surjective,

$$\begin{aligned}
 LC(Y, \sigma) = LC(f(X), \sigma) &= \bigwedge_{y \in f(x) \subseteq f(X)} \bigvee_{U' = f(U) \subseteq f(X)} [N_y^Y(U') \otimes [\Gamma(U', \sigma/U')]] \\
 &\geq \bigwedge_{y \in f(x) \subseteq f(X)} [N_{f(x)}^Y(f(U)) \otimes [\Gamma(f(U), \sigma/f(U))]] \\
 &\geq \bigwedge_{y \in f(x) \subseteq f(X)} [\tau(V') \otimes O(f) \otimes \Gamma_P(U, \tau/U) \otimes C_P(f)] \geq \lambda. \quad \square
 \end{aligned}$$

Theorem 4.7. *Let (X, τ) and (Y, σ) be two fuzzifying topological spaces and $f \in Y^X$ be surjective. Then $\models L_PC(X, \tau) \otimes I_P(f) \otimes O_P(f) \rightarrow L_PC(Y, \sigma)$.*

Proof. By Theorem 9 in [1], the proof is similar to the proof of Theorem 4.6. \square

Theorems 4.6 and 4.7 are a generalization of the following corollary.

Corollary 4.2. *Let (X, τ) and (Y, σ) be two topological spaces and $f : (X, \tau) \rightarrow (Y, \sigma)$ be surjective mapping. If f is a pre-continuous (resp. pre-irresolute), open (resp. pre-open) and X is locally strong compact, then Y is locally compact (resp. locally strong compact) space.*

Theorem 4.8. *Let $\{(X_s, \tau_s) : s \in S\}$ be a family of fuzzifying topological spaces, then*
 $= L_PC(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s) \rightarrow \forall s(s \in S \wedge L_PC(X_s, (\tau_P)_s) \wedge \exists T(T \sqsubset S \wedge \forall t(t \in S - T \wedge \Gamma_P(X_t, \tau_t)))$.

Proof. It suffices to show that

$$\begin{aligned}
 &L_PC(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s) \\
 &\leq \bigwedge_{s \in S} [L_PC(X_s, (\tau_P)_s) \wedge \bigvee_{T \sqsubset S} \bigwedge_{t \in S - T} \Gamma_P(X_t, \tau_t)].
 \end{aligned}$$

From Theorem 4.7 and Lemma 3.1 we have for any $t \in S$,

$$\begin{aligned}
 &L_PC(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s) \\
 &= [L_PC(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s) \otimes C_P(p_t) \otimes O_P(p_t)] \leq L_PC(X_t, \tau_t).
 \end{aligned}$$

So,

$$\bigwedge_{t \in S - T} L_PC(X_t, \tau_t) \geq L_PC(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s).$$

By Theorem 3.2 we have

$$\begin{aligned}
& \bigvee_{T \sqsubset S} \bigwedge_{t \in S-T} \Gamma_P(X_t, \tau_t) \\
& \geq [\bigvee_{U \subseteq \prod_{s \in S} X_s} \Gamma_P(U, \prod_{s \in S} (\tau_P)_s \setminus U) \otimes \bigvee_{X \subseteq \prod_{s \in S} X_s} N_x^{PX}(U)] \\
& \geq \bigvee_{U \subseteq \prod_{s \in S} X_s} \bigvee_{X \subseteq \prod_{s \in S} X_s} [\Gamma_P(U, \prod_{s \in S} (\tau_P)_s \setminus U) \otimes N_x^{PX}(U)] \\
& \geq \bigwedge_{X \subseteq \prod_{s \in S} X_s} \bigvee_{U \subseteq \prod_{s \in S} X_s} [\Gamma_P(U, \prod_{s \in S} (\tau_P)_s \setminus U) \otimes N_x^{PX}(U)] \\
& = L_P C(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s).
\end{aligned}$$

Therefore,

$$\begin{aligned}
& L_P C(\prod_{s \in S} X_s, \prod_{s \in S} (\tau_P)_s) \\
& \leq [\bigwedge_{t \in S-T} L_P C(X_t, \tau_t) \wedge \bigvee_{T \sqsubset S} \bigwedge_{t \in S-T} \Gamma_P(X_t, \tau_t)].
\end{aligned}$$

□

We can obtain the following corollary in crisp setting.

Corollary 4.3. *Let $\{X_\lambda : \lambda \in \Lambda\}$ be a family of nonempty topological spaces. If $\prod_{\lambda \in \Lambda} X_\lambda$ is locally strong compact, then each X_λ is locally strong compact and all but finitely many X_λ are strong compact*

Conclusion: The present paper investigates topological notions when these are planted into the framework of Ying's fuzzifying topological spaces (in semantic method of continuous valued-logic). It continue various investigations into fuzzy topology in a legitimate way and extend some fundamental results in general topology to fuzzifying topology. An important virtue of our approach (in which we follow Ying) is that we define topological notions as fuzzy predicates (by formulae of Łukasiewicz fuzzy logic) and prove the validity of fuzzy implications (or equivalences). Unlike the (more

wide-spread) style of defining notions in fuzzy mathematics as crisp predicates of fuzzy sets, fuzzy predicates of fuzzy sets provide a more genuine fuzzification; furthermore the theorems in the form of valid fuzzy implications are more general than the corresponding theorems on crisp predicates of fuzzy sets. The main contributions of the present paper are to give characterizations of fuzzifying strong compactness. Also, we define the concept of locally strong compactness of fuzzifying topological spaces and obtain some basic properties of such spaces. There are some problems for further study:

- (1) One obvious problem is: our results are derived in the Łukasiewicz continuous logic. It is possible to generalize them to more general logic setting, like residuated lattice-valued logic considered in [34-35].
- (2) What is the justification for fuzzifying locally strong compactness in the setting of $(2, L)$ topologies.
- (3) What is the justification for fuzzifying locally strong compactness in (M, L) -topologies etc.

References

- [1] S. A. Abd El-Baki and O. R. Sayed, Pre-irresoluteness and strong compactness in fuzzifying topology, *J. Egypt Math. Soc.*, Vol. 15 (1), pp. 41-56, 2007.
- [2] K. M. Abd El-Hakeim, F. M. Zeyada and O. R. Sayed, Pre-continuity and $D(c, p)$ -continuity in fuzzifying topology, *Fuzzy Sets and Systems*, Vol. 119, pp. 459-471, 2001.
- [3] K. M. Abd El-Hakeim, F. M. Zeyada and O. R. Sayed, Pre-separation axioms in fuzzifying topology, *Fuzzy Systems and Mathematics*, Vol. 17, pp. 29-37, 2003.
- [4] C. L. Chang, Fuzzy topological spaces, *J. Math. Anal. Appl.*, Vol. 24, pp. 182-190, 1968.
- [5] A. Dutta and B. C. Tripathy, On fuzzy $b\hat{I}$ open sets in fuzzy topological space, *Journal of Intelligent and Fuzzy Systems*, Vol. 32 (1), pp. 137-139, 2017.

- [6] J. A. Goguen, The fuzzy Tychonoff theorem, *J. Math. Anal. Appl.*, Vol. 43, pp. 182-190, 1973.
- [7] U. Höhle, Uppersemicontinuous fuzzy sets and applications, *J. Math. Anal. Appl.*, Vol. 78, pp. 659-673, 1980.
- [8] U. Höhle, Many Valued Topology and its Applications, *Kluwer Academic Publishers, Dordrecht*, 2001.
- [9] U. Höhle, S. E. Rodabaugh, Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory, in: Handbook of Fuzzy Sets Series, vol. 3, *Kluwer Academic Publishers, Dordrecht*, 1999.
- [10] U. Höhle, S. E. Rodabaugh, A. Šostak, (Eds.), Special Issue on Fuzzy Topology, *Fuzzy Sets and Systems*, Vol. 73, pp. 1-183, 1995.
- [11] U. Höhle, A. Šostak, Axiomatic foundations of fixed-basis fuzzy topology, in: U. Höhle, S. E. Rodabaugh, (Eds.), Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory, in: Handbook of Fuzzy Sets Series, Vol. 3, *Kluwer Academic Publishers, Dordrecht*, pp. 123-272, 1999.
- [12] J. L. Kelley, General Topology, *Van Nostrand, New York*, 1955.
- [13] T. Kubiak, On Fuzzy Topologies, Ph.D. Thesis, *Adam Mickiewicz University, Poznan, Poland*, 1985
- [14] Y. M. Liu, M. K. Luo, Fuzzy Topology, *World Scientific, Singapore*, 1998.
- [15] A. S. Mashhour, M. E. Abd El-Monsef and S. N. El-Deeb, On pre-continuous and weak pre-continuous mappings, *Proc. Math. Phs. Soc. Egypt*, Vol. 53, pp. 255-263, 1982.
- [16] A. S. Mashhour, M. E. Abd El-Monsef, I. A. Hasanein and T. Noiri, Strongly compact spaces, *Delta J. Sci.*, Vol. 8, pp. 30-64, 1984.
- [17] D. Qiu, Fuzzifying topological linear spaces, *Fuzzy Sets and Systems*, Vol. 147, pp. 249-272, 2004.
- [18] D. Qiu, Characterizations of fuzzy finite automata, *Fuzzy Sets and Systems*, Vol. 141 pp. 391-414, 2004.

- [19] G. C. Ray and B. C. Tripathy, On fuzzy δ^* -almost continuous and fuzzy δ^* -continuous functions in mixed fuzzy topological spaces, *Proyecciones J. Math.*, Vol. 39 (2), pp. 435-449, 2020.
- [20] S. E. Rodabaugh, Categorical foundations of variable-basis fuzzy topology, in: U. Höhle, S. E. Rodabaugh, (Eds.), *Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory*, in: *Handbook of Fuzzy Sets Series*, vol. 3, *Kluwer Academic Publishers, Dordrecht*, pp. 273-388, 1999.
- [21] J. B. Rosser, A. R. Turquette, *Many-Valued Logics*, *North-Holland, Amsterdam*, 1952.
- [22] D. J. Sarma and B. C. Tripathy, Fuzzy semi-pre quasi neighborhood structure, *Afrika Matematika*, Vol. 30 (1-2), pp. 217-221, 2019.
- [23] O. R. Sayed, On Fuzzifying Topological Spaces, Ph.D. Thesis, *Assiut University, Egypt*, 2002.
- [24] J. Shen, Locally compactness in fuzzifying topology, *J. Fuzzy Math.*, Vol. 2 (4), pp. 695-711, 1994.
- [25] B. C. Tripathy and G. C. Ray, Mixed fuzzy ideal topological spaces, *Applied Mathematics and Computations*, Vol. 220, pp. 602-607, 2013.
- [26] B. C. Tripathy and G. C. Ray, On δ -continuity in mixed fuzzy topological spaces, *Boletim da Sociedade Paranaense de Matematica*, Vol. 32 (2), pp. 175-187, 2014.
- [27] B. C. Tripathy and G. C. Ray, Weakly continuous functions on mixed fuzzy topological spaces, *Acta Scientiarum. Technology*, Vol. 36 (2), pp. 331-335, 2014 .
- [28] B. C. Tripathy and G. C. Ray, Fuzzy δ -I-continuity in mixed fuzzy ideal topological spaces, *Journal of Applied Analysis*, Vol. 24 (2), pp. 233-239, 2018.
- [29] G. J. Wang, *Theory of L-Fuzzy Topological Spaces*, *Shanxi Normal University Press, Xi an*, 1988 (in Chinese).
- [30] M. S. Ying, A new approach for fuzzy topology (I), *Fuzzy Sets and Systems*, Vol. 39, pp. 303-321, 1991.

- [31] M. S. Ying, A new approach for fuzzy topology (II), *Fuzzy Sets and Systems*, Vol. 47, pp. 221-23, 1992.
- [32] M. S. Ying, A new approach for fuzzy topology (III), *Fuzzy Sets and Systems*, Vol. 55, pp. 193-207, 1993.
- [33] M. S. Ying, Compactness in fuzzifying topology, *Fuzzy Sets and Systems*, Vol. 55, pp. 79-92, 1993.
- [34] M. S. Ying, Fuzzifying topology based on complete residuated lattice-valued logic (I), *Fuzzy Sets and Systems*, Vol. 56, pp. 337-373, 1993.
- [35] M. S. Ying, Fuzzy topology based on residuated lattice-valued logic, *Acta Mathematica Sinica*, Vol. 17, pp. 89-102, 2001.

O. R. Sayed

Department of Mathematics,
Faculty of Science,
Assiut University,
Assiut 71516,
Egypt
e-mail: o_sayed@aun.edu.eg
e-mail: o_r_sayed@yahoo.com
Corresponding author

and

Adem Kiliçman

Department of Mathematics,
University Putra Malaysia,
43400 UPM, Serdang, Selangor,
Malaysia
e-mail: akilic@upm.edu.my